

## SUPPLEMENTARY DATA

**Supplementary Table S1. Human genes associated with T1D: overlap with other autoimmune diseases and animal models of T1D.** Human association data were extracted from the Immunobase database ([www.immunobase.org](http://www.immunobase.org)) and from references (1-4). Light blue shading indicates that the same marker (single nucleotide polymorphism) is shared between the disease and T1D, while dark blue shading indicates that a different marker within or around the gene is associated with the disease, or in a few cases, that the gene has been associated without the exact marker being defined. Markers that are not associated with particular genes or that are associated with lincRNA were excluded from the table. The number of overlaps for each disease is shown after the disease abbreviation (out of a total of 64 loci listed for T1D). For rodent genetic studies, dark orange shading indicates that the gene(s) in the susceptibility region (Idd for NOD mice and Iddm for BB rats) has been implicated, while lighter shading simply denotes that the gene is within the susceptibility region, but has not been formally implicated. Mouse and rat association data were from t1dbase.org and from references (5; 6). At time of submission, however, the t1dbase.org database was no longer accessible, although a fraction of it has been archived with over 2,300 URLs accessible on [https://web.archive.org/web/\\*/t1dbase.org/\\*](https://web.archive.org/web/*/). Abbreviations: AA, Alopecia areata; AS, Ankylosing spondylitis; ATD, Autoimmune thyroid disease; CEL, Celiac disease; CRO, Crohn's disease; IBD, Inflammatory bowel disease; JIA, Juvenile Idiopathic Disease; MS, Multiple sclerosis; PBC, Primary biliary cirrhosis; PSC, Primary sclerosing cholangitis; PSO, Psoriasis; RA, Rheumatoid arthritis; SJO, Sjogren syndrome; SLE, Systemic lupus erythematosus; SSC, Systemic scleroderma; T1D, Type 1 diabetes, UC, Ulcerative colitis; VIT, Vitiligo.

SUPPLEMENTARY DATA

Loci	Main genes	T1D	CRO (32)	IBD (31)	UC (30)	MS (25)	RA (24)	CEL (23)	JIA (16)	SLE (15)	PBC (11)	VIT (11)	AA (10)	PSO (9)	PSC (8)	SSC (7)	AS (6)	ATD (5)	SJG (5)	NOD	BB
1p13.2	PTPN22, PHTF1																			Idd18.2	Iddm26
1p31.1	PGM1																				
1q31.2	RGS1																			Idd5.4	
1q32.1	IL10																			Idd5	
2p23.3	EFR3B																				
2q11.2	AFF3																			Idd26	
2q12.1	IL18RAP																				
2q24.2	IFIH1																				
2q32.3	STAT4																			Idd5	
2q33.2	CTLA4																			Idd5.1	
2q33.2	SLC11A1																			Idd5.2	
3p21.31	CCR5																				
4q27	IL2, IL21, ADAD1																			Idd3	Iddm32
5p13.2	IL7R																				
5p15	ERAP1																				
6q15	BACH2																				
6p21.3	PRSS16																				
MHC	HLA-DR, HLA-DQ																			Idd1	Iddm1
6q21	FYN																				
6q22.32	CENPW																				
6q23.3	TNFAIP3																			Idd23	
6q25.3	TAGAP																				
7p12.1	COBL																				
7p12.2	IKZF1																				
7p15.2	SKAP2																				Iddm14

## SUPPLEMENTARY DATA

## SUPPLEMENTARY DATA

**Supplementary Table S2. Human genes associated with T1D: known markers or SNPs, and function or potential function in APCs** (under parenthesis if function known in other cell types). Markers that are not associated with particular genes or that are associated with lincRNA were excluded. Genes that are known or expected to affect APC function are highlighted.

Loci	Main genes	SNPs	List of SNPs	Potential role in APCs	Refs
1p13.2	<b>PTPN22, PHTF1</b>	2	rs2476601, rs6679677	PTPN22 regulates type I interferon responses to TLR agonists	(7; 8)
1p31.1	PGM1	1	rs2269241	(unknown)	
1q31.2	RGS1	2	rs2209014, rs2816316	(T cells)	
1q32.1	<b>IL10</b>	2	rs3024493, rs3024505	IL-10 inhibits T cell effector response and supports regulatory T cell functions	(2)
2p23.3	EFR3B	1	rs478222	(unknown)	
2q11.2	AFF3	3	rs6740838, rs9653442, rs13415583	May play a role in monocytes	(9)
2q12.1	IL18RAP	1	rs917997	(IFNg secretion)	
2q24.2	<b>IFIH1</b>	4	rs1990760, rs2111485, rs35667974, rs72871627	IFIH1 regulates the production of type I interferon in response to viruses and RNA	(10)
2q32.3	<b>STAT4</b>	2	rs6744380, rs7574865	STAT4 is key to IL-12 signaling and is required for T1D development	(11; 12)
2q33.2	CTLA4	2	rs3087243, rs11571316	(T cells, Tregs)	
2q33.2	<b>SLC11A1</b>	1	rs3731685	Nramp1, endosomal protein (DCs, MΦs) enhances MHC-II expression and antigen presentation, silencing reduces disease incidence in NOD mice; also influences the IL-12/IL-10 balance	(13-18)
3p21.31	<b>CCR5</b>	4	rs333, rs11711054, rs17078977, rs113010081	DC / macrophage homing to inflamed sites Association controversial for certain variants	(19-23)
4q27	IL2, IL21, ADAD1	5	rs2069763, rs4505848, rs6827756, rs17388568, rs75793288	(T cells)	
5p13.2	IL7R	2	rs1445898, rs11954020	(lymphocytes)	
5p15	<b>ERAP1</b>	1	rs30187	Antigen processing: trimming of MHC class I peptides; effect on proinflammatory cytokines / chemokines	(24-26)
6q15	BACH2	3	rs597325, rs11755527, rs72928038	(lymphocytes)	
6p21.3	<b>PRSS16</b>	0	unconfirmed	Antigen processing (thymus)	(27-30)
MHC	<b>HLA-DR, HLA-DQ</b>	3	rs6916742, rs9272346, rs9268645	Antigen presentation: susceptibility conferred by HLA-DR3/DR4 and DQ8/DQ2, protection from DQ6	(31; 32)
6q21	FYN	1	rs11964650	(unknown)	
6q22.32	CENPW	3	rs9388489, rs1538171, rs9375435	(unknown)	
6q23.3	<b>TNFAIP3</b>	3	rs1878658, rs2327832, rs6920220	Encodes A20, a negative regulator of NF-κB and TLR signaling; expression of A20 is reduced in LADA	(33; 34)
6q25.3	<b>TAGAP</b>	2	rs212402, rs1738074	Macrophages from Tagap-deficient mice produce less type I interferons, TNF-α, IL-6 and inflammatory chemokines in response to HSV infection	(19; 35)
7p12.1	COBL	1	rs4948088	(actin regulation)	
7p12.2	<b>IKZF1</b>	3	rs62447205, rs10272724, rs201847125	IKAROS, encoded by IKZF1, regulates the development and differentiation of mDCs and pDCs	(36; 37)
7p15.2	<b>SKAP2</b>	2	rs7804356, rs12533947	Required for MΦ migration and chemotaxis	(38)
8p23.1	<b>CTSB</b>	1	rs1296023	Protease contributing to antigen processing	(39)
9p24.2	GLIS3	3	rs6476839, rs7020673, rs10758593	(pancreatic beta cells)	
10p11.22	<b>NRP1, ITGB1</b>	3	rs722988, rs1557150, rs2666236	NRP1: expressed at high levels on pDCs; it is unclear whether its ligation can potentiate tolerogenic functions as seen with Tregs; ITGB1-encoded CD29 is an adhesion molecule expressed on some myeloid cells and involved in the proliferation of β-cells	(39-41)
10p15.1	IL2RA, RBM17	6	rs2104286, rs7090530, rs10795791, rs12251307, rs41295121, rs61839660	(T cells, Tregs)	
10p15.1	PRKCQ	2	rs2236380, rs11258747	(T cells, NF-κB signaling downstream of TCR)	
10q23.31	RNLS	2	rs10509540, rs12416116	May play a role in monocytes	(9)

## SUPPLEMENTARY DATA

11p15.5	<b>INS, IGF2</b>	5	rs689, rs3741208, rs7111341, rs7928968, rs72853903	INS: autoantigen; level of expression in thymic epithelial cells influences selection of insulin-reactive T cells; mutations can favor binding of INS DRiP on MHC.	(42-44)
11q13.1	<b>BAD</b>	1	rs694739	(apoptosis)	
12p13.31	<b>CD69</b>	3	rs917911, rs4763879, rs10492166	Regulates tissue retention and egress, including that of DCs	(2; 45)
12q13.2	<b>ERBB3, IKZF4</b>	5	rs705704, rs705705, rs705708, rs2292239, rs11171739	ERBB3: expressed in maturing DCs and monocytes, with higher expression being more protective; (IKZF4: role in T cells and Tregs.)	(46)
12q13.2	<b>DGKA, RAB5B</b>	1	rs11171710	(T cells)	
12q13.13	<b>ITGB7</b>	1	rs11170466	The protein associates with CD103 (ITGA6) or CD49d (ITGA4) to regulate leukocyte homing	(39)
12q14.1	<b>CYP27B1</b>	2	rs4760341, rs10877012	Responsible for the production of bioactive vitamin D, which potentiates the tolerogenic function of DCs	(47)
12q24.12-13	<b>SH2B3, NAA25</b>	3	rs653178, rs3184504, rs17696736	Lnk (encoded by SH2B3) regulates the number of DCs and their response to GM-CSF	(19; 48; 49)
13q22	<b>LMO7</b>	1	rs539514	(unknown)	
13q32.3	<b>GPR183</b>	2	rs6491500, rs9585056	Stimulation of GPR183-encoded EBI2 regulates cell migration and positioning of macrophages and DCs	(9; 50; 51)
14q24.1	<b>ZFP36L1</b>	1	rs1465788	Regulated monocyte / macrophage differentiation	(52)
14q32.2	<b>DLK1</b>	2	rs941576, rs56994090	Expressed on mucosal DCs	
15q14	<b>RASGRP1</b>	4	rs7171171, rs12908309, rs17574546, rs72727394	(lymphocytes)	
15q25.1	<b>CTSH</b>	3	rs3825932, rs12148472, rs34593439	Protease contributing to antigen processing	(53)
16p11.2	<b>IL27, NUPR1</b>	3	rs151234, rs4788084, rs9924471	IL-27: induces IL-10+ Tr1 cells and also potentiate Foxp3+ Tregs	(2; 54; 55)
16p13.13	<b>CLEC16A, DEXI</b>	3	rs193778, rs12708716, rs12927355	CLEC16A regulates autophagy and antigen processing in APCs; DEXI may play a role in monocytes	(9; 56-58)
16q23.1	<b>CTRB1</b>	1	rs7202877	(pancreas)	
17p13.2	<b>UBE2G1</b>	1	rs9906760	(unknown)	
17q12	<b>GSDMB, ORMDL3</b>	3	rs2290400, rs12150079, rs12453507	(asthma ORMDL3)	
17q21.2	<b>CCR7, SMARCE1</b>	1	rs757411	CCR7: enable DC homing to lymph nodes; SMARCE1: may play a role in monocytes	(9; 49)
17q21.31	<b>MAP3K14</b>	1	rs17759555	Encodes the NF- $\kappa$ B-inducing kinase (NIK), which regulates the NF- $\kappa$ B pathway and cross-priming by DCs	(39; 59)
18p11.21	<b>PTPN2</b>	3	rs1893217, rs2542151, rs12971201	Besides a major role in T cells, it also regulates M-CSF receptor signaling and macrophage differentiation	(60)
18q22.2	<b>CD226</b>	3	rs763361, rs1615504, rs1790575	CD226 cross-linking enhances MHC-II, CD80 and CD86 on CD8 $\alpha$ + DCs, but not CD8 $\alpha$ - DCs	(9; 61)
19p13.2	<b>TYK2</b>	3	rs1051738, rs12720356, rs34536443	TYK2 regulates the production of IL-12 by DCs, as well as the response to IL-12	(12)
19q13.32	<b>PRKD2</b>	2	rs425105, rs60652743	(mostly implicated in tumor invasion); may play a role in monocytes	(9)
19q13.33	<b>FUT2</b>	3	rs516246, rs601338, rs602662	(intestinal epithelial cells)	
20p13	<b>SIRPG</b>	1	rs2281808	(T cells)	
21q22.3	<b>UBASH3A</b>	2	rs11203202, rs11203203	(T cells: inhibition of NF- $\kappa$ B signaling under TCR)	
22q12.2	<b>HORMAD2</b>	1	rs5753037	(unknown)	
22q12.3-13.1	<b>C1QTNF6, RAC2</b>	2	rs229533, rs229541	C1QTNF6: regulation of IL-10 production; RAC2: encodes a small GTPase that controls phagosome pH and cross-presentation	(62; 63)
22q12.3	<b>IL2RB</b>	1	rs229526	(T cells)	
Xp22.2	<b>TLR8</b>	1	rs5979785	Response of DCs and other APCs to ssRNA	(64)
Xq28	<b>GAB3</b>	1	rs2664170	Encodes an adaptor protein that binds GRB2 and SHP2 tyrosine phosphatase and regulated monocyte / macrophage differentiation; functions downstream of M-CSF receptor	(65)

## SUPPLEMENTARY DATA

**Supplementary Table S3. Selected rodent genes associated with T1D with a known or putative role in APCs.** The table shows a limited selection of relevant genes from the reviews by Driver et al. (5) and Wallis et al. (6). Genes present in Idd regions are NOD mice, while those in Iddm regions are from diabetes-prone BB rats.

Main genes	Region	Potential role in APCs	Refs
<i>B2m</i>	Idd13.1, Iddm27	Component of MHC-I, $\beta$ 2m variation can influence the selection of peptides for MHC-I binding and susceptibility to disease	(66; 67)
<i>C1qtnf6</i>	Iddm34	Enhances IL-10 production by M $\Phi$ s, see Table S2	(63)
<i>Cd101</i>	Idd10/Idd18.4	Crosslinking induces IL-10 secretion by DCs	(5; 68)
<i>Csf1</i>	Idd18.3	Encodes M-CSF, a growth factor for M $\Phi$ s	Table S6
<i>Csf2</i>	Idd4.3	Encodes GM-CSF, a growth factor for DCs	Table S6
<i>Iapp</i>	Idd6.2	Autoantigen, for tolerance induction? Accumulation in islets contributes to inflammation and macrophage activation	(69)
<i>Il1a / Il1b</i>	Idd13, Iddm27	Proinflammatory cytokines	(5; 6)
<i>Il10</i>	Idd5	Immunoregulatory cytokine produced by DCs	Table S2
<i>Il27</i>	Iddm10	Immunoregulatory cytokine produced by DCs	Table S2
<i>Ins1 / Ins2</i>	Iddm25	Autoantigen, for tolerance induction	Table S2
<i>MHC-II</i>	Idd1, Iddm1	Permissible haplotype I-A <sup>g7</sup> (NOD mouse) and RT1 <sup>u</sup> (BB rat)	
<i>Nfkb1</i>	Iddm33	Encodes subunit 1 of NF- $\kappa$ B	Table S11
<i>Sirpa</i>	Idd13.2	NOD variant on BM-DCs binds more strongly to CD47	(70)
<i>Slc11a1</i>	Idd5.2	Influences antigen presentation and cytokine production	Table S2
<i>Vdr</i>	Iddm34	Encodes the vitamin D receptor	(6)
Unidentified	Idd4	Controls overproduction of IL-12 by M $\Phi$ s	(71)
Unidentified	Idd10/17/18	Controls overproduction of IL-12 by DCs arrested in maturing phase	(72)

## SUPPLEMENTARY DATA

**Supplementary Table S4. Altered number of DC and monocyte populations in T1D.** The table is divided in two sections: A) blood DCs and monocytes and B) in vitro differentiated DCs from monocytes. Abbreviations: mDCs: myeloid DCs; pDCs: plasmacytoid DCs; MoDCs: monocyte-derived DCs; ctrl: control; NO: new onset T1D; LT: long-term T1D; AAb: autoantibodies; A: adults; C: children; YA: young adults;  $\nearrow$  increased in T1D;  $\searrow$  decreased in T1D;  $\equiv$  no change.

(\*) Trend for more mDCs and less pDCs in T1D (not significant). Lin- cells typically exclude lymphocytes, NK cells and monocytes.

	APC	Phenotype	Subject	Age group	Frequency / cell count	Refs
A	Blood DCs	Lin- HLA-DR+	T1D vs ctrl	A	Frequency $\equiv$ Cell count $\equiv$	(73)
	Blood DCs	Lin- HLA-DR+	T1D At-risk, NO vs LT, ctrl	All	Frequency $\nearrow$	(74)
	Blood mDCs	Lin- HLA-DR+ CD11c+ CD123lo	T1D vs ctrl	A	Frequency $\equiv$ (*)	(73)
	Blood mDCs	CD1c+ CD19-	T1D vs ctrl	C/YA	Frequency $\equiv$	(75)
	Blood mDCs	CD1c+ CD19-	AAb+ vs ctrl	All	Frequency $\equiv$	(76)
	Blood mDCs	Lin- HLA-DR+ CD11c+ CD123-	T1D At-risk, NO, LT vs ctrl	YA	Cell count $\equiv$	(77)
	Blood mDCs	Lin- HLA-DR+ CD11c+ CD123-	T1D At-risk, NO, LT vs ctrl	All	Frequency $\searrow$	(74)
	Blood mDCs	HLA-DR+ CD11c+ CD1c+	T1D vs ctrl	C	Frequency $\searrow$ Cell count $\searrow$	(78)
	Blood mDCs	Lin- HLA-DR+ CD11c+	T1D NO, LT vs ctrl	C	Frequency $\searrow$ Cell count $\searrow$	(79)
	Blood mDCs	cDC1: Lin- CD1c+	T1D NO vs ctrl	All	Frequency $\searrow$	(80)
	Blood mDCs	cDC1: Lin- CD1c+	T1D LT vs ctrl	All	Frequency $\equiv$	(80)
	Blood pDCs	Lin- HLA-DR+ CD11c- CD123lo	T1D vs ctrl	A	Frequency $\equiv$ (*)	(73)
	Blood pDCs	CD304+	T1D vs ctrl	C/YA	Frequency $\equiv$	(75)
	Blood pDCs	CD304+	AAb+ vs ctrl	All	Frequency $\equiv$	(76)
	Blood pDCs	Lin- HLA-DR+ CD11c- CD123+	T1D At-risk, NO, LT vs ctrl	All	Frequency $\nearrow$	(74)
	Blood pDCs	HLA-DR+ CD123+ CD303+	T1D vs ctrl	C	Frequency $\searrow$ Cell count $\searrow$	(78)
	Blood pDCs	Lin- HLA-DR+ CD123+ CD304+	T1D NO, LT vs ctrl	C	Frequency $\searrow$ Cell count $\searrow$	(79)
	Blood pDCs	Lin- CD303+ CD304+	T1D NO, LT vs ctrl	All	Frequency $\searrow$ Cell count $\searrow$	(80)
	Blood pDCs	Lin- HLA-DR+ CD11c- CD123+	T1D At-risk, NO, LT vs ctrl	YA	Frequency $\nearrow$ Cell count $\equiv$	(77)
B	Blood monocytes	CD14+	T1D vs ctrl	C/YA	Frequency $\equiv$	(75)
	Blood monocytes	CD14+	AAb+ vs ctrl	All	Frequency $\equiv$	(76)
	Blood monocytes	HLA-DR+ CD14+ CD19-	T1D NO vs ctrl	C	Frequency $\nearrow$ Cell count $\nearrow$	(79)
	Blood monocytes	HLA-DR+ CD14+ CD19-	T1D LT vs ctrl	C	Frequency $\equiv$ Cell count $\searrow$	(79)
	MoDCs	N/A	T1D vs ctrl	All	Yield $\searrow$	(81)
	MoDCs	HLA-DR+ CD14-	AAb+ vs ctrl	All	Yield $\searrow$	(82)
	MoDCs	HLA-DR+ CD11c+	At-risk, NO vs ctrl	C	Yield $\searrow$	(83)

## SUPPLEMENTARY DATA

**Supplementary Table S5. Altered frequency and yield of DCs from NOD mice compared to other mouse strains.** The table is divided in two sections: A) freshly isolated DCs and B) in vitro differentiated DCs from bone marrow. The cytokines listed are those used for in vitro differentiation of the DCs. “Not seen in” means that the reduced frequency of CD8 $\alpha$ + DCs observed in the spleen was not seen in the indicated tissues. Abbreviations: BM-DCs: bone marrow-derived DCs; PLN: pancreatic lymph node;  $\nearrow$  increased in NOD;  $\searrow$  decreased in NOD. (\*) Phenotype: CD11c<sup>low</sup> mPDCA (CD317)+.

APC		Mouse strains	Frequency / yield	Notes	Refs
A	Splenic CD8 $\alpha$ + DCs	NOD vs B6, B10.DR	Frequency $\searrow$		(84)
	Splenic CD8 $\alpha$ + DCs	NOD vs NOR, B6, BALB/c	Frequency $\searrow$	Not seen in PLNs	(85)
	Splenic CD8 $\alpha$ + DCs	NOD vs NOR, B6	Frequency $\searrow$	Not seen in thymus	(86)
	Splenic CD8 $\alpha$ + DCs	NOD.H-2k vs B10.H-2k	Frequency $\searrow$	In favor of CD8 $\alpha$ - DCs; Idd1-independent	(87)
	Splenic pDCs (*)	NOD (young vs diabetic)	Frequency $\searrow$		(88)
	Splenic DCs	NOD vs CBA, DBA/2	Frequency $\searrow$		(89)
	Splenic DCs	NOD vs DBA/2	In vitro yield $\searrow$	FLT3L	(89)
	PLN merocytic DCs	NOD vs B6, BALB/c	Frequency $\nearrow$	CD11c+ CD8 $\alpha$ - CD11b-/low	(90)
B	Liver B220+ DCs	NOD vs NOR, BALB/c (females)	Yield $\searrow$	Defective in vitro propagation	(91)
	BM-DCs	NOD vs B6, CBA, DBA/2	Yield $\searrow$	GM-CSF/IL-4	(89)
	BM-DCs	NOD vs B6, DBA/2	Yield $\searrow$	GM-CSF/IL-4; bias toward granulocytes	(92)
	BM-DCs	NOD vs B6	Yield $\searrow$	GM-CSF $\pm$ IL-4	(93)
	BM-DCs	NOD vs B10.DR	Yield $\searrow$	GM-CSF/IL-4	(94)
	BM-DCs	NOD vs B10.DR, B6, DBA/2	Yield $\searrow$	GM-CSF $\pm$ IL-4	(95)
	BM-DCs	NOD vs NOR, B6, BALB/c	Yield $\searrow$	GM-CSF; bias toward MΦ-like cells	(96)
	BM-DCs	NOD.H-2k vs B10.H-2k	Yield $\searrow$	GM-CSF/IL-4; Idd1-independent	(97)
	BM-DCs	NOD vs NOR, B6, BALB/c	Yield $\nearrow$	GM-CSF/IL-4	(98)

## SUPPLEMENTARY DATA

**Supplementary Table S6. Role of growth factors in APC alterations and defects in NOD mice.**

Cytokine	Role / effect	Refs
FLT3L	Treatment with FLT3L corrects deficiency in CD8α+ DC numbers in NOD mice	(84; 86)
	Treatment with FLT3L expands pDC precursors and restore their function in NOD mice	(99)
GM-CSF	Greater number of granulocyte-macrophage colony-forming cells (NOD vs B6)	(100)
	Reduced response of myeloid progenitors to GM-CSF (NOD vs BALB/c)	(101)
	Enhanced proliferation of pancreatic DC precursors to GM-CSF (NOD vs B6)	(102)
	Over-production of GM-CSF linked to <i>Csf2</i> polymorphism (Idd4.3) in monocytes and MΦs (NOD vs B6)	(103)
	Persistent STAT5 stimulation by GM-CSF that is impervious to regulation by IL-10 (NOD vs B6; T1D / high-risk patients vs healthy controls)	(104)
	Increased GM-CSF production and STAT5 phosphorylation in monocytes from T1D patients and at-risk subjects	(105)
M-CSF	Reduced yield of MΦs from bone marrow cells with M-CSF (NOD vs NON and congenics)	(106)

## SUPPLEMENTARY DATA

**Supplementary Table S7. Distribution of extra-pancreatic expression of major  $\beta$ -cell antigens.** The cell-specific and tissue-specific expression of autoantigens (outside the pancreas) is reported according to published reports and gene expression databases including the Immunological Genome Project (ImmGen, [www.ImmGen.org](http://www.ImmGen.org)), BioGPS ([www.biogps.org](http://www.biogps.org)) and the Genotype-Tissue Expression Project (GTEx, [www.gtexportal.org](http://www.gtexportal.org)). Although not a  $\beta$ -cell antigen, GAD67 is listed because several epitopes are identical (cross-reactive) between GAD65 and GAD67.

Islet autoantigen	Extra-pancreatic tissue	Notes, other sources	Refs
Proinsulin ( <i>INS</i> )	Thymus	In mTECs, AIRE-dependent	(107-109)
	Lymph nodes (fibroblastic reticular cells)	In FRCs (mice), nodes in mice and humans	(110-113)
	Blood myeloid cells	Humans (monocytes, DCs)	(114)
	Peripheral DCs	Human and mouse	(115-118)
	Liver, bone marrow, adipose tissue	Under hyperglycemic conditions (mice)	(119)
Chromogranin A ( <i>CHGA</i> )	Adrenal and pituitary glands, stomach	BioGPS (mouse) and GTEx (human)	
	Adrenal, pituitary and thyroid glands	BioGPS (human)	
	mTECs	ImmGen (mouse)	
IA-2 ( <i>PTPRN</i> )	Brain, pituitary and adrenal glands	BioGPS (mouse) and GTEx (human)	
	Brain / pineal gland	BioGPS (human)	
	Thymus, spleen	Splice variant missing exon 13 only	(120)
	mTECs	ImmGen (mouse)	(109)
IAPP ( <i>IAPP</i> )	mTECs	ImmGen (mouse)	
	pDCs (?)	BioGPS (mouse)	
IGRP ( <i>G6PC2</i> )	mTECs (low)	ImmGen (mouse)	
	Thymus, spleen	Splice variants differ from pancreas	(121; 122)
	Testis?	GTEx (human)	
GAD65 ( <i>GAD2</i> )	Brain (NOT expressed in mTECs)	BioGPS and GTEx (human)	(109)
GAD67 ( <i>GAD1</i> )	Brain	BioGPS (mouse, human) and GTEx (human)	
	mTECs and lymph node stromal cells	ImmGen (mouse)	(109; 111; 123)
ZnT8 ( <i>SLC30A8</i> )	Testis	GTEx (human)	

## SUPPLEMENTARY DATA

**Supplementary Table S8. Alterations in antigen processing and peptide loading in T1D.**

Process	Gene product	Association with T1D and relevant functions	Refs
Antigen processing	CLEC16A	Polymorphism associated with T1D	(9; 56)
		Regulates autophagy in humans	(57)
		Knockdown of CLEC16A in NOD mice protects from T1D by impacting autophagy in mTECs and T cell selection	(58)
	CTSB	Polymorphism associated with T1D	(39)
		Regulates antigen processing, autophagy and lysosomal dynamics	(124; 125)
	CTSH	Polymorphism associated with T1D	(53)
		Regulates antigen processing	(125)
		Polymorphism associated with lower insulin production in $\beta$ cells (a role in mTECs and other cells?)	(126)
	ERAP1	Polymorphism associated with T1D	(24)
		Regulates processing of antigens to MHC class I and influencing recognition by CD8+ T cells; associated with multiple autoimmune diseases	(25)
		Effect on proinflammatory cytokine expression	(26)
	PRSS16	Polymorphism associated with T1D	(27; 28)
		PRSS16 knock-out NOD mice were protected from disease, in part due to a more effective deletion of certain diabetogenic CD4+ T cells in the thymus	(29; 30)
Peptide loading	CLIP	Elevated levels of I-A $^{g7}$ -associated CLIP on splenic B cells (NOD vs other strains)	(127)
		Increased % of CLIP+ monocytes in T1D patients (vs control and T2D patients)	(128)
		Increased density of empty HLA-DR on B cells from T1D patients (vs control twins)	(129)
		Poor HLA-DM editing leads to persistence of CLIP	(130)
		HLA-DQ2 and DQ8 are more resistant to CLIP release by HLA-DM	(131)
	INS-DRiP	Polymorphism in the 3'UTR of INS leads to better binding of the insulin defective ribosomal product (DRiP) to HLA-DQ8/DQ2 molecules. T1D patients who carry both susceptible DQ8/DQ2 and INS-DRiP alleles have T cell responses to this neoantigen, while subjects carrying the protective DQ6 or INS-DRiP allele alone do not show any response.	(44)
	HIP	Hybrid insulin peptides: fusion products between the C-terminus of InsC-derived peptide and the N-terminus of a peptide from another $\beta$ -cell antigen (chromogranin A or IAPP) fit the I-A $^{g7}$ MHC particularly well. Such peptides were also identified in T1D patients as HLA-DQ8 or DQ8/DQ2-restricted.	(132; 133)

## SUPPLEMENTARY DATA

**Supplementary Table S9. Altered expression of MHC and costimulatory molecules in APCs from NOD mice compared to other mouse strains.** The table is divided in two sections: A) MHC, and B) Costimulatory molecules CD40, CD80 and CD86. Abbreviations: pDCs: Plasmacytoid DCs; LPS: lipopolysaccharides; PIC: poly I:C; ↑ increased in NOD; ↓ decreased in NOD; ≡ no change. Data refer to mean fluorescence intensity, except (#) based on % cells positive for the cytokine and (\*) based on mRNA levels. (\*\*) pDCs defined as SiglecH<sup>+</sup> BST2<sup>+</sup> B220<sup>+</sup>.

	APC	Mouse strains	Stimulation	MHC and costimulation	Refs
MHC molecules	Splenic DCs (ex vivo)	NOD vs DBA/2	None	MHC-II ↓	(89)
	Splenic DCs (cultured)	NOD vs DBA/2	None	MHC-II ↓	(89)
	BM-DCs	NOD vs B6	None	MHC-II ↓	(96)
	BM-DCs	NOD vs B6	None	MHC-II ↓	(93)
	BM-DCs	NOD vs DBA/2	None	MHC-II ↓	(89)
	BM-DCs	NOD vs DBA/2	None	MHC-II ↓	(92)
	BM-DCs	NOD vs CBA	None, LPS/IFN-γ	MHC-II ↓	(134)
	BM-DCs	NOD vs B10.DR	None	MHC-II ≡	(94)
	BM-DCs	NOD.H-2k vs B10.H-2k	None	MHC-II ↓ (#)	(97)
	BM-DC	NOD vs B6	LPS, CD40L	MHC-II ↓	(72)
	Pre-DCs	NOD vs NOR	CpG	MHC-II ↓	(99)
	Splenic MΦs	NOD vs NOR, BALB/c	None, LPS	MHC-I ↑ MHC-II ≡	(135)
	Peritoneal MΦs	NOD vs NOR, B6	IFN-γ	MHC-I ↓	(106)
Costimulatory molecules	Splenic CD8α+ DCs	NOD vs B6	None	CD40 ↑	(136)
	Splenic CD8α+ DCs	NOD vs B6	None	CD80, CD86 ≡ CD40 ↑	(137)
	Splenic CD8α+ DCs	NOD vs B6	CpG	CD80, CD86 ↓ CD40 ↑	(137)
	Splenic CD8α+ DCs	NOD vs NOR, BALB/c	None, LPS, PIC	CD40, CD80, CD86 ≡	(85)
	Splenic CD8α- DCs	NOD vs NOR, BALB/c	None, LPS, PIC	CD40, CD80, CD86 ≡	(85)
	Splenic CD8α- DCs	NOD vs B6	None, CpG	CD80, CD86 ≡ CD40 ↑	(137)
	Splenic DCs (ex vivo)	NOD vs DBA/2	None	CD40 ↑ CD80, CD86 ≡	(89)
	Splenic DCs (in vitro)	NOD vs DBA/2	None	CD80 ↓ (#) CD40, CD86 ≡	(89)
	Splenic pDCs (**)	NOD vs B6	None	CD40, CD80, CD86 ≡	(137)
	Splenic pDCs (**)	NOD vs B6	CpG	CD40 ≡ CD80, CD86 ↓	(137)
	Pancreatic CD8α- DCs	NOD vs B6	None	CD40, CD86 ↓	(138)
	BM-DCs	NOD vs NOR, B6, BALB/c	None	CD40, CD80, CD86 ≡	(96)
	BM-DCs	NOD vs BALB/c	None	CD40, CD80, CD86 ≡	(139)
	BM-DCs	NOD vs BALB/c, B6	None	CD40, CD80, CD86 ≡	(140)
	BM-DCs	NOD vs BALB/c, B6	None	CD80 ↓ CD40, CD86 ≡	(141)
	BM-DCs	NOD vs B6	None	CD40, CD80, CD86 ↓	(93)
	BM-DCs	NOD vs CBA	None	CD40, CD80, CD86 ↓	(134)
	BM-DCs	NOD vs DBA/2	None	CD40, CD86 ↓	(92)
	BM-DCs	NOD vs B10.DR, B6	None	CD40, CD80, CD86 ↑	(95)
	BM-DCs	NOD vs B10.DR	None	CD40 ↑	(94)
	BM-DCs	NOD vs BALB/c, B6	None	CD40, CD80 ↑	(142)
	BM-DCs	NOD, NOR vs BALB/c, B6	None	CD40, CD80, CD86 ↑	(139)
	BM-DCs	NOD vs DBA/2	None	CD80 ↓ (#) CD40, CD86 ≡	(89)
	BM-DCs	NOD vs NON	LPS	CD80, CD86 ↓ (*)	(143)
	BM-DCs	NOD vs CBA	LPS/IFN-γ	CD40 ≡ CD80, CD86 ↓	(134)
	BM-DCs	NOD vs BALB/c	LPS	CD40, CD80, CD86 ≡	(139)
	BM-DCs	NOD vs BALB/c, B6	LPS	CD40, CD80, CD86 ≡	(140)
	BM-DCs	NOD vs B6	LPS, CD40L	CD86 ↓	(72)
	BM-DCs	NOD vs BALB/c, B6	LPS/IFN-γ	CD40,80,86 ≡	(141)
	BM-DCs	NOD vs BALB/c, B6	With T cells	CD80,86 ↑	(141)
	Pre-DCs	NOD vs NOR	CpG	CD80,86 ↓	(99)
	Splenic MΦs	NOD vs NOR, BALB/c	None	CD40, CD80, CD86 ≡	(135)
	Splenic MΦs	NOD vs NOR, BALB/c	LPS	CD40, CD80 ≡ CD86 ↑	(135)
	B cells	NOD vs BALB/c, B6, B6g7	None	CD40, CD80 ↑	(142)

## SUPPLEMENTARY DATA

**Supplementary Table S10. Altered expression of MHC and costimulatory molecules in APCs from T1D patients compared to various controls.** The table is divided in two sections: A) MHC, and B) Costimulatory molecules CD40, CD80 and CD86. Abbreviations: MoDCs: monocyte-derived DCs; mDCs: myeloid DCs; pDCs: plasmacytoid DCs; NO: new onset; ctrl: control; FDR: first degree relative; LPS: lipopolysaccharides;  $\nearrow$  increased in T1D;  $\searrow$  decreased in T1D;  $\equiv$  no change. Data refer to mean fluorescence intensity, except (#) based on % cells positive for the cytokine and (\*) based on mRNA levels.

	APC	Subject	Stimulation	MHC and costimulation	Refs
MHC molecules	Blood DCs	T1D NO vs ctrl (children)	None	HLA-DR $\equiv$	(79)
	Blood mDCs	T1D vs ctrl	None	HLA-DR $\nearrow$	(74)
	Blood mDCs	T1D vs ctrl	None	HLA-DR $\nearrow$	(77)
	Blood pDCs	T1D vs ctrl	None	HLA-DR $\nearrow$	(74)
	Blood pDCs	T1D vs ctrl	None	HLA-DR $\nearrow$	(77)
	MoDCs	T1D, At-Risk vs ctrl (children)	None	HLA-DR $\searrow$ (#)	(83)
	MoDCs	T1D vs ctrl (children)	None, LPS	HLA-DR $\equiv$	(144)
	MoDCs	T1D vs ctrl, T2D, RA	LPS	HLA-DR $\equiv$	(145)
Costimulatory molecules	PBMCs	T1D NO vs ctrl (children)	None	CD80 $\nearrow$ CD40, 86 $\equiv$ (*)	(146)
	PBMCs	T1D NO vs ctrl (adults)	None	CD86 $\nearrow$ CD40, 80 $\equiv$ (*)	(146)
	Blood DCs	T1D NO vs ctrl (children)	None	CD40, CD86 $\equiv$	(79)
	Blood mDCs	T1D vs ctrl	None	CD40, CD80, CD86 $\equiv$	(77)
	Blood mDCs	T1D vs ctrl	None	CD40, CD80, CD86 $\equiv$ (#)	(73)
	Blood pDCs	T1D vs ctrl	None	CD40, CD80, CD86 $\equiv$	(77)
	Blood pDCs	T1D vs ctrl	None	CD40, CD80, CD86 $\equiv$ (#)	(73)
	MoDCs	T1D vs ctrl, T2D	None	CD80 $\equiv$ CD86 $\nearrow$	(147)
	MoDCs	T1D vs ctrl (children)	None	CD80 $\searrow$ (#)	(83)
	MoDCs	At-Risk FDR vs ctrl	None	CD80, CD86 $\searrow$ (#)	(82)
	MoDCs	T1D vs ctrl	None	CD40, CD80, CD86 $\equiv$	(148)
	MoDCs	T1D vs ctrl (children)	None	CD80, CD86 $\equiv$	(144)
	MoDCs	T1D vs ctrl, T2D	None, TNF/IL-1/IL-6	CD40, CD80, CD86 $\equiv$	(149)
	MoDCs	T1D vs ctrl (children)	LPS	CD80, CD86 $\searrow$	(144)
	MoDCs	T1D vs ctrl, T2D, RA	LPS	CD40 $\searrow$	(145)
	MoDCs	T1D vs ctrl, T2D, RA	LPS	CD80, CD86 $\equiv$	(145)
	MoDCs	T1D vs ctrl	TNF- $\alpha$	CD80 $\nearrow$ CD40, CD86 $\equiv$	(148)

## SUPPLEMENTARY DATA

**Supplementary Table S11. Evidence of the implication of the NF-κB pathway in the abnormal maturation of DCs and MΦs in T1D.**

Species	Role / effect	Refs
NOD mice	Hyperactive IκB kinase and NF-κB in splenic DCs and BM-DCs (NOD vs B6 and BALB/c)	(140)
	Excessive NF-κB activity enhances costimulation and cytokine secretion (NOD vs NOR, BALB/c)	(139)
	Overexpression of IL-12 by MΦs is due to abnormal regulation by NF-κB and linked to Idd4	(71; 150)
	Excessive NF-κB activity in BM-DCs and B cells (NOD vs BALB/c, B6)	(142)
	Lower expression of p65/p50/p52 and RelB in BM-DCs (NOD vs NON)	(143)
BB rats	Polymorphism in the <i>NFKB1</i> gene (Iddm33)	(6)
T1D patients	Excessive NF-κB activity in monocytes and monocyte-derived DCs from T1D patients	(145)
	Susceptibility gene <i>MAP3K14</i> encodes NIK, a positive regulator of NF-κB pathway	Table S2
	Susceptibility gene <i>TNFAIP3</i> encodes A20, a negative regulator of NF-κB pathway	Table S2

## SUPPLEMENTARY DATA

**Supplementary Table S12A. Altered cytokine expression in APCs from NOD mice compared to other mouse strains.** The table is divided in four sections: IL-10, IL-12, proinflammatory cytokines, and Type I interferons (T1). Abbreviations: LPS: lipopolysaccharides; PIC: poly I:C; PGN: peptidoglycan; SAC: Staph. aureus Cowan extract; ↑ increased in NOD; ↓ decreased in NOD; ≡ no change. Data refer to MFI or amount secreted, except (\*) indicating mRNA levels.

	<b>APC</b>	<b>Mouse strains</b>	<b>Stimulation</b>	<b>Cytokines</b>	<b>Refs</b>
IL-10	Splenic CD8α+ DCs	NOD vs NOR, BALB/c	PIC	IL-10 ≡	(85)
	Splenic CD8α+ DCs	NOD vs NOR, BALB/c	CD40L/LPS	IL-10 ↑	(85)
	Splenic CD8α- DCs	NOD vs NOR, BALB/c	PIC, CD40L/LPS	IL-10 ≡	(85)
	Pancreatic CD8α+ DCs	NOD vs B6	None	IL-10 ↓ (*)	(151)
	Pancreatic CD8α- DCs	NOD vs B6	None	IL-10 ↓ (*)	(138)
	Pancreatic CD8α- DCs	NOD vs B6	LPS	IL-10 ≡	(138)
	Pancreatic CD8α- DCs	NOD vs B6	LPS	IL-10 ↑ (*)	(138)
	BM-DCs	NOD vs B6	LPS	IL-10 ↓	(72)
	Macrophages	NOD vs B6, B10	LPS	IL-10 ↓	(152)
	Macrophages	NOD vs NOR	LPS	IL-10 ↑	(152)
IL-12	Splenic CD8α+ DCs	NOD vs B6	CpG	IL-12p40 ↓	(86)
	Splenic CD8α+ DCs	NOD vs NOR, BALB/c	PIC, CD40L/LPS	IL-12 ↓	(85)
	Splenic CD8α+ DCs	NOD vs B6	LPS, PIC, SAC	IL-12p40 ↓	(84)
	Splenic CD8α- DCs	NOD vs NOR, BALB/c	PIC, CD40L/LPS	IL-12 ≡	(85)
	Splenic DCs	NOD vs NOR, BALB/c	PIC	IL-12 ↓ IL-12p40 ↓	(85)
	Splenic DCs	NOD vs NOR	CD40L/LPS	IL-12 ↓ IL-12p40 ↓	(85)
	Splenic DCs	NOD vs B6, B10.BR	SAC, TNF-α	IL-12p40 ↓	(84)
	Splenic DCs	NOD vs B6	CpG	IL-12	(137)
	Pancreatic CD8α- DCs	NOD vs B6	None	IL-12b ↓ (*)	(138)
	Pancreatic CD8α- DCs	NOD vs B6	LPS	IL-12 ≡	(138)
	Pancreatic CD8α- DCs	NOD vs B6	LPS	IL-12a ↑ (*)	(138)
	BM-DCs	NOD vs BALB/c, B6	CD40L, LPS	IL-12 ↑	(140)
	BM-DCs	NOD vs NOR, BALB/c	CD40L, LPS	IL-12 ↑	(139)
	BM-DCs	NOD vs DBA/2	LPS/CD40L	IL-12 ↑	(89)
	BM-DCs	NOD vs B6	LPS	IL-12 ↑	(72)
	BM-DCs	NOD vs B6	LPS/T cells	IL-12 ↑	(141)
	BM-DCs	NOD vs B10.DR	None	IL-12p40 ↑	(94)
	BM-DCs	NOD vs B10.DR	LPS	IL-12p40 ≡	(94)
	BM-DCs	NOD vs NOR	LPS	IL-12a ↓ (*)	(143)
	BM-DCs	NOD vs B6, B10.DR	CD40L or LPS/IFN-γ	IL-12 ↓	(95)
	BM-DCs	NOD vs B6	LPS/IFN-γ	IL-12 ↓	(93)
	Macrophages	NOD vs NOR	LPS	IL-12 ↑	(85)
	Macrophages	NOD vs B6	LPS	IL-12 ↑	(153)
	Macrophages	NOD vs B6, B10, NOR	LPS	IL-12 ↑	(152)
	Macrophages	NOD vs A/J, B6, BALB/c	LPS	IL-12 ↑ IL-12p40 ↑	(150)
	Macrophages	NOD vs A/J, B6, BALB/c	LPS	IL-12p40 ↑	(71)
	Macrophages	NOD vs NOR, BALB/c	LPS, TNF-α, CD40L	IL-12 ↑	(135)
	Macrophages	NOD vs NOR, BALB/c	LPS	IL-12p40 ↑ (*)	(135)

## SUPPLEMENTARY DATA

**Supplementary Table S12B. Altered cytokine expression in APCs from NOD mice compared to other mouse strains.** The table is divided in four sections: IL-10, IL-12, proinflammatory cytokines, and Type I interferons (T1). Abbreviations: LPS: lipopolysaccharides; PIC: poly I:C; PGN: peptidoglycan; SAC: Staph. aureus Cowan extract; ↑ increased in NOD; ↓ decreased in NOD; ≡ no change. Data refer to MFI or amount secreted, except (\*) indicating mRNA levels.

APC	Mouse strains	Stimulation	Cytokines	Refs
Pro-inflammatory cytokines	Splenic CD8α+ DCs	NOD vs B6	LPS, PIC, SAC	IL-6 ≡ (84)
	Splenic CD8α- DCs	NOD vs B6	LPS, PIC, SAC	IL-6 ↑ (84)
	Splenic DCs	NOD vs NOR	PIC	TNF-α ≡ (85)
	Splenic DCs	NOD vs B6	CpG	TNF-α ↑ (137)
	Pancreatic CD8α- DCs	NOD vs B6	LPS	IL-6 ↑ TNF-α ≡ (138)
	Pancreatic CD8α- DCs	NOD vs B6	LPS	IL-1b, IL-6, TNF ↑ (*) (138)
	BM-DCs	NOD vs NON	LPS	IL-1a, IL-1b, IL-6 ↓ (*) (143)
	BM-DCs	NOD vs B6, B10.DR	LPS/IFN-γ	IL-6 ≡ (95)
	Pre-DCs	NOD vs NOR	CpG	IL-6, TNF-α ↓ (99)
	Macrophages	NOD vs B6	PGN	TNF-α ↓ (153)
	Macrophages	NOD vs NOR, BALB/c	LPS	IL-1α, TNF-α ↑ (135)
	Macrophages	NOD vs NOR, BALB/c	CD40L	TNF-α ↑ IL-1α ≡ (135)
	Macrophages	NOD vs NOR	LPS	IL-1, TNF-α ↓ IL-6 ≡ (152)
	Macrophages	NOD vs B6, B10	LPS	IL-1, TNF-α ≡ IL-6 ↑ (152)
IFN-I	Splenic DCs	NOD vs B6	CpG	IFN-α ↑ IFN-β ↑ (137)
	Splenic pDCs	NOD (young vs diabetic)	None (basal)	IFN-α ↓ (88)
	BM-DC (Flt3L)	NOD vs B6g7	CpG	IFN-α ↑ (137)

## SUPPLEMENTARY DATA

**Supplementary Table S13. Altered cytokine expression in APCs from T1D patients compared to various controls.** The table is divided in four sections: A) IL-10, B) IL-12, C) proinflammatory cytokines, and D) Type I interferons. Abbreviations: MoDCs and MoMΦs: monocyte-derived DCs and macrophages; mDCs: myeloid DCs; LPS: lipopolysaccharides; PIC: poly I:C; PC4: Pam3CSK4; ctrl: control; AAb: autoantibody; FDR: first degree relative; ↑ increased in T1D; ↓ decreased in T1D; ≡ no change. Data refer to mean fluorescence intensity or amount secreted, except (#) based on % cells positive for the cytokine and (\*) based on mRNA levels.

	<b>APC</b>	<b>Subjects</b>	<b>Stimulation</b>	<b>Cytokines</b>	<b>Refs</b>
IL-10	Blood DCs	New onset T1D (children)	PIC	IL-10 ≡ (#)	(79)
	MoDCs	T1D vs ctrl, T2D, RA	LPS	IL-10 ≡	(145)
	MoDCs	T1D or At-Risk vs ctrl	IL-1β	IL-10 =	(83)
	MoDCs	T1D vs ctrl (children)	LPS	IL-10 ↑	(144)
	Blood monocytes	T1D vs ctrl, T2D (adults)	LPS	IL-10 ↑	(147)
	Blood monocytes	New onset T1D (children)	PIC	IL-10 ≡ (#)	(79)
	Blood monocytes	T1D vs ctrl, T2D	IFN-γ	IL-10 ≡ (*)	(154)
	B cells	T1D vs ctrl	IL-21	IL-10 ≡ (#)	(155)
	B cells	T1D, LADA vs ctrl, T2D	None	B10 cells ↓ (#)	(156)
	B cells	T1D vs ctrl	Anti-IgM	IL-10 ↓ (#) IL-10 ↓	(157)
	B cells	T1D vs ctrl	PC4, LPS, CpG	IL-10 ↓	(157)
IL-12	Blood DCs	T1D vs ctrl, T2D	CD40L, LPS	IL-12 ↓	(73)
	Blood DCs	New onset T1D (children)	PIC	IL-12 ≡ (#)	(79)
	MoDCs	T1D vs ctrl (children)	LPS	IL-12 ↓	(144)
	MoDCs	T1D vs ctrl	IL-1β	IL-12 ≡	(83)
	MoDCs	At-Risk vs ctrl	IL-1β	IL-12 ↓	(83)
	Blood monocytes	New onset T1D (children)	PIC	IL-12 ≡ (#)	(79)
	Blood monocytes	T1D vs ctrl, T2D	LPS	IL-12 ≡	(147)
Proinflammatory cytokines	Blood monocytes	T1D vs ctrl, T2D	IFN-γ	IL-12 ↑ (*)	(154)
	Blood DCs	New onset T1D (children)	PIC	IL-6, TNF-α ≡ (#)	(79)
	Blood mDCs	New onset T1D (children)	LPS, PIC, R848	IL-1β ≡ (#)	(75)
	Blood mDCs	New onset T1D (children)	LPS	IL-6 ↓ (#)	(75)
	Blood mDCs	New onset T1D (children)	PIC, R848	IL-6 ≡ (#)	(75)
	Blood mDCs	AAb+ vs AAb-	None, LPS, PIC	IL-1β, IL-6 ≡ (#)	(76)
	Blood mDCs	AAb+ vs AAb-	R848	IL-1β ↑ (#) IL-6 ≡ (#)	(76)
	MoDCs	T1D vs ctrl, T2D, RA	LPS	IL-1β, IL-6, TNF-α ≡	(145)
	MoDCs	T1D vs ctrl	TNF-α	IL-6 ↑	(148)
	MoDCs	T1D or At-Risk vs ctrl	IL-1β	TNF-α ↓ IL-6 ≡	(83)
	Blood monocytes	New onset T1D (children)	None, LPS	IL-1β ↑ (#) IL-6 ≡ (#)	(75)
	Blood monocytes	New onset T1D (children)	PIC, R848	IL-1β, IL-6 ≡ (#)	(75)
	Blood monocytes	AAb+ vs AAb-	None, LPS	IL-1β ↑ (#) IL-6 ≡ (#)	(76)
	Blood monocytes	AAb+ vs AAb-	PIC, R848	IL-1β ↑ (#) IL-6 ↓ (#)	(76)
	Blood monocytes	New onset T1D (children)	PIC	IL-6 ≡ (#)	(79)
	Blood monocytes	New onset T1D (children)	PIC	TNF-α ↑ (#)	(79)
	Blood monocytes	T1D vs ctrl	None, LPS, PC4	IL-1β ↑ TNF-α ↑	(158)
	Blood monocytes	T1D vs ctrl, T2D	LPS	IL-6 ↑ IL-1β, TNF-α ≡	(147)
IFN-1	MoMΦs	DQ8/DQ2 T1D vs ctrls	LPS	IL-1β, IL-6, TNF-α ↑	(159)
	B cells	T1D vs ctrl	Anti-IgM	TNF-α ↑ (#) TNF-α ↓	(157)
	Blood DCs	T1D, T2D vs ctrl	CD40L	IFN-α ↓	(73)
	Blood pDCs	New onset T1D (children)	R848 (TLR7/8)	IFN-α ↑ (#)	(75)
	Blood pDCs	FDR vs T1D, ctrl	CpG	IFN-α ↑	(160)
PBMCs	PBMCs	T1D patients	Flu virus	IFN-α ↑	(74)
	PBMCs	T1D patients	Flu virus	IFN-α ↑	(77)

## SUPPLEMENTARY DATA

**Supplementary Table S14. Other mechanisms of tolerance that are altered in APCs in T1D.** The column H/M/R indicates whether the data refers to humans, mice or rats. Abbreviations: IDO: indoleamine 2,3-dioxygenase; MS: multiple sclerosis.

Mechanism	Role / effect	H/M/R	Refs
Galectin-1	Inhibitor of T cell proliferation and Th1/Th17 responses Monocytes produce less galectin-1 in T1D patients compared to controls	H	(161)
IDO	Inhibitor of T cell proliferation and effector T cell response in favor of Treg responses	H/M/R	(162)
	Contribute to the protective role of pDCs in NOD mice	M	(163)
	DCs from young female NOD mice have an impaired induction of IDO in response to IFN- $\gamma$ , relative to male NOD mice or B6 mice	M	(164)
	Defects due to inhibited Stat1 signaling, corrected by peroxynitrite inhibitor or a Stat1-independent inducer (chorionic gonadotrophin)	M	(164; 165)
	Defective IDO induction by IFN- $\gamma$ in NOD mice extends to dermal fibroblasts	M	(166)
Vitamin D	Regulator of the tolerogenic function of DCs	H/M/R	(167)
	<i>CYP27B1</i> , encoding a critical enzyme for the generation of vitamin D, is associated with T1D and MS	H	Table S1 Table S2
	<i>CYP27B1</i> polymorphism can lead to reduced expression in DCs (MS)	H	(168)
	<i>CYP27B1</i> polymorphism leads to reduced expression in PBMCs (T1D)	H	(169)
	One <i>CYP27B1</i> polymorphism is associated with lower Treg frequency (T1D)	H	(170)
	<i>VDR</i> , encoding the vitamin D receptor, is associated with T1D in BB rats	R	Table S3

## SUPPLEMENTARY DATA

**Supplementary Table S15. Defects in adhesion and homing of APCs.** The table is divided in two sections: A) cell adhesion, and B) chemotaxis. The column H/M indicates whether the data refers to humans (H) or mice (M). Abbreviations: AAb: autoantibodies; MoDCs: monocyte-derived DCs; mDCs: myeloid DCs.

Molecule	Role / effect	H/M	Refs
Adhesion molecules	ICAM-1	H	(147)
	ITGB7	M	(171-173)
	<i>ITGB7</i> is associated with human T1D	H	Table S2
	<i>ITGB1</i> is associated with human T1D	H	Table S2
	<i>ITGB1</i> may also be required for $\beta$ -cells to proliferate	M	(41)
	ITGA4/B7 and ITGB1 can bind fibronectin	H/M	
	BM-DCs have increased adhesion to fibronectin in vitro (NOD vs B6, BALB/c)	M	(174)
SLAM	Reduced SLAM expression in NOD mice leads to diminished homotypic interaction between mDCs and NKT cells (NOD vs B6) and to defective secretion of IL-4 and IL-10 by NKT cells, which play a regulatory role in T1D	M	(175; 176)
SIRP $\alpha$	The protein encoded by the NOD allele of <i>Sirpa</i> on DCs binds more strongly to CD47, which results in enhanced T cell proliferation	M	(70)
Chemotaxis	Impaired migration toward inflamed tissue sites (NOD vs B6, BALB/c)	M	(177)
Chemokines and receptors	CCR2	M	(138)
	<i>CCL2</i> (CCR2 ligand) recruits tolerogenic CD11c+ CD11b+ DCs to islets (NOD)	M	(178)
	CCR2-deficient NOD mice have lower incidence of disease diabetes	M	(179)
	Reduced on mDCs and pDCs from children with T1D (vs controls)	H	(78)
	Blood levels of <i>CCL2</i> (CCR2 ligand) are reduced in subjects with multiple AAb.	H	(180)
	CCR2 association with T1D is controversial	H	(21-23)
	CCR5	M	(138)
	Involved in the recruitment of leukocytes, including DCs, to inflamed islets	M	(181; 182)
	CCR5-deficient NOD mice have accelerated diabetes	M	(179)
	Blockade of <i>CCL5</i> (one of CCR5 ligands) ameliorates diabetes	M	(181)
	Levels of <i>CCL3</i> and <i>CCL4</i> (other CCR5 ligands) are increased in blood of subjects with multiple AAb	H	(180)
	<i>CCR5</i> is associated with human T1D	H	Table S2
CCR7	<i>CCR7</i> is associated with human T1D	H	Table S2
	Defective migration of mature NOD BM-DCs toward <i>CCL19</i> (a CCR7 ligand)	M	(174)
CXCL12	Association of <i>CXCL12</i> with early onset of T1D	H	(183; 184)
	Increased expression of CXCL12 in NOD pancreas recruits pDCs (NOD vs NOR, B6)	M	(185)

## SUPPLEMENTARY DATA

**Supplementary Table S16. Preclinical studies featuring exogenous APCs ± antigen provision.** Abbreviations: BM-DCs bone marrow-derived DCs; IV intravenous; IP intraperitoneal; ID intradermal; MPLA monophosphoryl lipid A.

Study details	Outcome	Refs
<b>Feili-Hariri et al. 1999.</b> BM-DCs (4 days with GM-CSF +/- IL-4), unpulsed or pulsed with (HSP60(437-460), GAD65(509-528), GAD65(524-543), 60 µg/ml); 4-8x10 <sup>5</sup> DCs injected IV in NOD mice (three weekly injections at 5-7 weeks)	DCs were protective with no added benefit from antigen provision	(186)
<b>Papaccio et al. 2000.</b> Splenic DCs, unpulsed or pulsed with human gamma globulin (100 µg/ml); 3x10 <sup>5</sup> DCs injected IV in NOD mice (one injection at 8 wks)	Protection required antigen pulsing (irrelevant antigen)	(187)
<b>Krueger et al. 2003.</b> Spleen-derived DC line, unpulsed or pulsed with insulin (400 µg/ml); 4x10 <sup>5</sup> APCs injected IP in cyclophosphamide-treated NOD mice (one injection at 8 weeks)	Protection with insulin-DCs, no protection with OVA as antigen	(188)
<b>Lo et al. 2006.</b> BM-DCs (6 days with GM-CSF and IL-4), unpulsed or pulsed with 3 µM InsB(9-23) or InsC19-A3 or GAD65(78-97) peptide; 1x10 <sup>5</sup> DCs injected in footpad of NOD mice (three injections from 9 to 11 weeks)	Only DCs with GAD65 peptide had a significant protection	(189)
<b>Marin-Gallen et al. 2010.</b> BM-DCs (7 days with GM-CSF), co-cultured with apoptotic bodies from NIT-1 cells (beta cell line); 1x10 <sup>6</sup> DCs injected IP in RIP-IFN-β NOD mice (one single injection at 12-14 days of age)	Only DCs loaded with beta cell apoptotic cells protected, compared to unpulsed or pulsed with apoptotic cells from non-beta cells.	(190)
<b>Haase et al. 2010.</b> BM-DCs (8 days with GM-CSF and IL-10, 1.5% mouse serum), pulsed with InsB(9-23) and InsB(15-23) at 10 µg/ml; 1x10 <sup>6</sup> DCs injected IP in NOD mice (weekly injections from 5 to 12 weeks of age)	Only DCs pulsed with insulin peptides were protective	(191)
<b>Gibson et al. 2015.</b> BM-DCs (6 days with GM-CSF, IL-4 and vitamin D3), matured with LPS for 24h, pulsed with InsC19-A3 (10 µg/ml); 2x10 <sup>5</sup> DCs injected ID in humanized HLA-DR4 Tg mice (two injections two weeks apart)	Reduced proliferation and IFNγ, increased IL-10 in ex vivo recall response	(192)
<b>Lo et al. 2018.</b> BM-DCs (5-6 days with GM-CSF and IL-4), used immature and pulsed with 3 µM InsB(9-23), GAD65(78-97) or GAD65(260-279) peptide; 1x10 <sup>5</sup> DCs injected in footpad of NOD mice (three weekly injections starting at 9 weeks of age).	Pulsing with certain subdominant and ignored epitopes conferred better protection than unpulsed	(193)
<b>Funda et al. 2018.</b> BM-DCs (7 days with GM-CSF and IL-4, 10% FBS or serum-free medium; vitamin D2 and dexamethasone added on day 6), unpulsed or pulsed with GAD65 (1-2 µg/ml) or OVA (1 µg/ml), and activated with MPLA for another day; 3x10 <sup>6</sup> DCs injected IP in NOD.SCID mice along with 5x10 <sup>6</sup> splenocytes from diabetic NOD mice. Also 3x10 <sup>6</sup> DCs injected IP in NOD mice (one single injection at 4 weeks of age).	Protective response with unpulsed DCs, which was lost when DCs were pulsed with GAD65 or OVA	(194)

## SUPPLEMENTARY DATA

### REFERENCES

1. Pociot F, Akolkar B, Concannon P, Erlich HA, Julier C, Morahan G, Nierras CR, Todd JA, Rich SS, Nerup J: Genetics of type 1 diabetes: what's next? *Diabetes* 2010;59:1561-1571
2. Barrett JC, Clayton DG, Concannon P, Akolkar B, Cooper JD, Erlich HA, Julier C, Morahan G, Nerup J, Nierras C, Plagnol V, Pociot F, Schuilenburg H, Smyth DJ, Stevens H, Todd JA, Walker NM, Rich SS, Type 1 Diabetes Genetics C: Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nature genetics* 2009;41:703-707
3. Bradfield JP, Qu HQ, Wang K, Zhang H, Sleiman PM, Kim CE, Mentch FD, Qiu H, Glessner JT, Thomas KA, Frackelton EC, Chiavacci RM, Imielinski M, Monos DS, Pandey R, Bakay M, Grant SF, Polychronakos C, Hakonarson H: A genome-wide meta-analysis of six type 1 diabetes cohorts identifies multiple associated loci. *PLoS genetics* 2011;7:e1002293
4. Pociot F, Lernmark A: Genetic risk factors for type 1 diabetes. *Lancet* 2016;387:2331-2339
5. Driver JP, Chen YG, Mathews CE: Comparative genetics: synergizing human and NOD mouse studies for identifying genetic causation of type 1 diabetes. *The review of diabetic studies* : RDS 2012;9:169-187
6. Wallis RH, Wang K, Marandi L, Hsieh E, Ning T, Chao GY, Sarmiento J, Paterson AD, Poussier P: Type 1 diabetes in the BB rat: a polygenic disease. *Diabetes* 2009;58:1007-1017
7. Wang Y, Shaked I, Stanford SM, Zhou W, Curtsinger JM, Mikulski Z, Shaheen ZR, Cheng G, Sawatzke K, Campbell AM, Auger JL, Bilgic H, Shoyama FM, Schmeling DO, Balfour HH, Jr., Hasegawa K, Chan AC, Corbett JA, Binstadt BA, Mescher MF, Ley K, Bottini N, Peterson EJ: The autoimmunity-associated gene PTPN22 potentiates toll-like receptor-driven, type 1 interferon-dependent immunity. *Immunity* 2013;39:111-122
8. Wang Y, Ewart D, Crabtree JN, Yamamoto A, Baechler EC, Fazeli P, Peterson EJ: PTPN22 Variant R620W Is Associated With Reduced Toll-like Receptor 7-Induced Type I Interferon in Systemic Lupus Erythematosus. *Arthritis Rheumatol* 2015;67:2403-2414
9. Wallace C, Rotival M, Cooper JD, Rice CM, Yang JH, McNeill M, Smyth DJ, Niblett D, Cambien F, Cardiogenics C, Tiret L, Todd JA, Clayton DG, Blankenberg S: Statistical colocalization of monocyte gene expression and genetic risk variants for type 1 diabetes. *Hum Mol Genet* 2012;21:2815-2824
10. Gorman JA, Hundhausen C, Errett JS, Stone AE, Allenspach EJ, Ge Y, Arkatkar T, Clough C, Dai X, Khim S, Pestal K, Liggitt D, Cerosaletti K, Stetson DB, James RG, Oukka M, Concannon P, Gale M, Jr., Buckner JH, Rawlings DJ: The A946T variant of the RNA sensor IFIH1 mediates an interferon program that limits viral infection but increases the risk for autoimmunity. *Nat Immunol* 2017;18:744-752
11. Yang Z, Chen M, Ellett JD, Fialkow LB, Carter JD, McDuffie M, Nadler JL: Autoimmune diabetes is blocked in Stat4-deficient mice. *J Autoimmun* 2004;22:191-200
12. Tokumasa N, Suto A, Kagami S, Furuta S, Hirose K, Watanabe N, Saito Y, Shimoda K, Iwamoto I, Nakajima H: Expression of Tyk2 in dendritic cells is required for IL-12, IL-23, and IFN-gamma production and the induction of Th1 cell differentiation. *Blood* 2007;110:553-560
13. Stober CB, Brode S, White JK, Popoff JF, Blackwell JM: Slc11a1, formerly Nramp1, is expressed in dendritic cells and influences major histocompatibility complex class II expression and antigen-presenting cell function. *Infection and immunity* 2007;75:5059-5067
14. Esposito L, Hill NJ, Pritchard LE, Cucca F, Muxworthy C, Merriman ME, Wilson A, Julier C, Delepine M, Tuomilehto J, Tuomilehto-Wolf E, Ionesco-Tirgoviste C, Nistico L, Buzzetti R, Pozzilli P, Ferrari M, Bosi E, Pociot F, Nerup J, Bain SC, Todd JA: Genetic analysis of chromosome 2 in type 1 diabetes: analysis of putative loci IDDM7, IDDM12, and IDDM13 and candidate genes NRAMP1 and IA-2 and the interleukin-1 gene cluster. *IMDIAB Group. Diabetes* 1998;47:1797-1799

## SUPPLEMENTARY DATA

15. Bassuny WM, Ihara K, Matsuura N, Ahmed S, Kohno H, Kuromaru R, Miyako K, Hara T: Association study of the NRAMP1 gene promoter polymorphism and early-onset type 1 diabetes. *Immunogenetics* 2002;54:282-285
16. Wicker LS, Chamberlain G, Hunter K, Rainbow D, Howlett S, Tiffen P, Clark J, Gonzalez-Munoz A, Cumiskey AM, Rosa RL, Howson JM, Smink LJ, Kingsnorth A, Lyons PA, Gregory S, Rogers J, Todd JA, Peterson LB: Fine mapping, gene content, comparative sequencing, and expression analyses support Ctla4 and Nramp1 as candidates for Idd5.1 and Idd5.2 in the nonobese diabetic mouse. *J Immunol* 2004;173:164-173
17. Kissler S, Stern P, Takahashi K, Hunter K, Peterson LB, Wicker LS: In vivo RNA interference demonstrates a role for Nramp1 in modifying susceptibility to type 1 diabetes. *Nature genetics* 2006;38:479-483
18. Dai YD, Marrero IG, Gros P, Zaghouani H, Wicker LS, Sercarz EE: Slc11a1 enhances the autoimmune diabetogenic T-cell response by altering processing and presentation of pancreatic islet antigens. *Diabetes* 2009;58:156-164
19. Smyth DJ, Plagnol V, Walker NM, Cooper JD, Downes K, Yang JH, Howson JM, Stevens H, McManus R, Wijmenga C, Heap GA, Dubois PC, Clayton DG, Hunt KA, van Heel DA, Todd JA: Shared and distinct genetic variants in type 1 diabetes and celiac disease. *The New England journal of medicine* 2008;359:2767-2777
20. Buhler MM, Craig M, Donaghue KC, Badhwar P, Willis J, Manolios N, Tait BD, Silink M, Bennetts BH, Stewart GJ: CCR5 genotyping in an Australian and New Zealand type 1 diabetes cohort. *Autoimmunity* 2002;35:457-461
21. Szalai C, Csaszar A, Czinner A, Szabo T, Panczel P, Madacsy L, Falus A: Chemokine receptor CCR2 and CCR5 polymorphisms in children with insulin-dependent diabetes mellitus. *Pediatr Res* 1999;46:82-84
22. Gambelunghe G, Ghaderi M, Brozzetti A, Del Sindaco P, Gharizadeh B, Nyren P, Hjelmstrom P, Nikitina-Zake L, Sanjeevi CB, Falorni A, Umbria Type 1 Diabetes R: Lack of association of CCR2-64I and CCR5-Delta 32 with type 1 diabetes and latent autoimmune diabetes in adults. *Human immunology* 2003;64:629-632
23. Yang B, Houlberg K, Millward A, Demaine A: Polymorphisms of chemokine and chemokine receptor genes in Type 1 diabetes mellitus and its complications. *Cytokine* 2004;26:114-121
24. Fung EY, Smyth DJ, Howson JM, Cooper JD, Walker NM, Stevens H, Wicker LS, Todd JA: Analysis of 17 autoimmune disease-associated variants in type 1 diabetes identifies 6q23/TNFAIP3 as a susceptibility locus. *Genes Immun* 2009;10:188-191
25. Fierabracci A, Milillo A, Locatelli F, Fruci D: The putative role of endoplasmic reticulum aminopeptidases in autoimmunity: insights from genomic-wide association studies. *Autoimmun Rev* 2012;12:281-288
26. Aldhamen YA, Pepelyayeva Y, Rastall DP, Seregin SS, Zervoudi E, Koumantou D, Aylsworth CF, Quiroga D, Godbehere S, Georgiadis D, Stratikos E, Amalfitano A: Autoimmune disease-associated variants of extracellular endoplasmic reticulum aminopeptidase 1 induce altered innate immune responses by human immune cells. *Journal of innate immunity* 2015;7:275-289
27. Lie BA, Akselsen HE, Bowlus CL, Gruen JR, Thorsby E, Undlien DE: Polymorphisms in the gene encoding thymus-specific serine protease in the extended HLA complex: a potential candidate gene for autoimmune and HLA-associated diseases. *Genes Immun* 2002;3:306-312
28. Lie BA, Viken MK, Akselsen HE, Flam ST, Pociot F, Nerup J, Kockum I, Cambon-Thomsen A, Thorsby E, Undlien DE: Association analysis in type 1 diabetes of the PRSS16 gene encoding a thymus-specific serine protease. *Human immunology* 2007;68:592-598
29. Viret C, Leung-Theung-Long S, Serre L, Lamare C, Vignali DA, Malissen B, Carrier A, Guerder S: Thymus-specific serine protease controls autoreactive CD4 T cell development and autoimmune diabetes in mice. *J Clin Invest* 2011;121:1810-1821

## SUPPLEMENTARY DATA

30. Viret C, Mahiddine K, Baker RL, Haskins K, Guerder S: The T Cell Repertoire-Diversifying Enzyme TSSP Contributes to Thymic Selection of Diabetogenic CD4 T Cell Specificities Reactive to ChgA and IAPP Autoantigens. *J Immunol* 2015;195:1964-1973
31. Noble JA: Immunogenetics of type 1 diabetes: A comprehensive review. *J Autoimmun* 2015;64:101-112
32. Pugliese A, Gianani R, Moromisato R, Awdeh ZL, Alper CA, Erlich HA, Jackson RA, Eisenbarth GS: HLA-DQB1\*0602 is associated with dominant protection from diabetes even among islet cell antibody-positive first-degree relatives of patients with IDDM. *Diabetes* 1995;44:608-613
33. Zhang M, Peng LL, Wang Y, Wang JS, Liu J, Liu MM, Hu J, Song B, Yang HB: Roles of A20 in autoimmune diseases. *Immunol Res* 2016;64:337-344
34. Cheng L, Zhang D, Jiang Y, Deng W, Wu Q, Jiang X, Chen B: Decreased A20 mRNA and protein expression in peripheral blood mononuclear cells in patients with type 2 diabetes and latent autoimmune diabetes in adults. *Diabetes Res Clin Pract* 2014;106:611-616
35. Liao J, Jijon HB, Kim IR, Goel G, Doan A, Sokol H, Bauer H, Herrmann BG, Lassen KG, Xavier RJ: An image-based genetic assay identifies genes in T1D susceptibility loci controlling cellular antiviral immunity in mouse. *PloS one* 2014;9:e108777
36. Wu L, Nichogiannopoulou A, Shortman K, Georgopoulos K: Cell-autonomous defects in dendritic cell populations of Ikaros mutant mice point to a developmental relationship with the lymphoid lineage. *Immunity* 1997;7:483-492
37. Allman D, Dalod M, Asselin-Paturel C, Delale T, Robbins SH, Trinchieri G, Biron CA, Kastner P, Chan S: Ikaros is required for plasmacytoid dendritic cell differentiation. *Blood* 2006;108:4025-4034
38. Alenghat FJ, Baca QJ, Rubin NT, Pao LI, Matozaki T, Lowell CA, Golan DE, Neel BG, Swanson KD: Macrophages require Skap2 and Sirpalpha for integrin-stimulated cytoskeletal rearrangement. *J Cell Sci* 2012;125:5535-5545
39. Evangelou M, Smyth DJ, Fortune MD, Burren OS, Walker NM, Guo H, Onengut-Gumuscu S, Chen WM, Concannon P, Rich SS, Todd JA, Wallace C: A method for gene-based pathway analysis using genomewide association study summary statistics reveals nine new type 1 diabetes associations. *Genet Epidemiol* 2014;38:661-670
40. Delgoffe GM, Woo SR, Turnis ME, Gravano DM, Guy C, Overacre AE, Bettini ML, Vogel P, Finkelstein D, Bonnevier J, Workman CJ, Vignali DA: Stability and function of regulatory T cells is maintained by a neuropilin-1-semaphorin-4a axis. *Nature* 2013;501:252-256
41. Diaferia GR, Jimenez-Caliani AJ, Ranjitkar P, Yang W, Hardiman G, Rhodes CJ, Crisa L, Cirulli V: beta1 integrin is a crucial regulator of pancreatic beta-cell expansion. *Development* 2013;140:3360-3372
42. Pugliese A, Zeller M, Fernandez A, Jr., Zalcberg LJ, Bartlett RJ, Ricordi C, Pietropaolo M, Eisenbarth GS, Bennett ST, Patel DD: The insulin gene is transcribed in the human thymus and transcription levels correlated with allelic variation at the INS VNTR-IDDM2 susceptibility locus for type 1 diabetes. *Nature genetics* 1997;15:293-297
43. Vafiadis P, Bennett ST, Todd JA, Nadeau J, Grabs R, Goodyer CG, Wickramasinghe S, Colle E, Polychronakos C: Insulin expression in human thymus is modulated by INS VNTR alleles at the IDDM2 locus. *Nature genetics* 1997;15:289-292
44. Kracht MJ, van Lummel M, Nikolic T, Joosten AM, Laban S, van der Slik AR, van Veelen PA, Carlotti F, de Koning EJ, Hoeben RC, Zaldumbide A, Roep BO: Autoimmunity against a defective ribosomal insulin gene product in type 1 diabetes. *Nat Med* 2017;23:501-507
45. Lamana A, Martin P, de la Fuente H, Martinez-Munoz L, Cruz-Adalia A, Ramirez-Huesca M, Escribano C, Gollmer K, Mellado M, Stein JV, Rodriguez-Fernandez JL, Sanchez-Madrid F, del Hoyo GM: CD69 modulates sphingosine-1-phosphate-induced migration of skin dendritic cells. *J Invest Dermatol* 2011;131:1503-1512

## SUPPLEMENTARY DATA

46. Wang H, Jin Y, Reddy MV, Podolsky R, Liu S, Yang P, Bode B, Reed JC, Steed RD, Anderson SW, Steed L, Hopkins D, Huang Y, She JX: Genetically dependent ERBB3 expression modulates antigen presenting cell function and type 1 diabetes risk. *PloS one* 2010;5:e11789
47. Bailey R, Cooper JD, Zeitels L, Smyth DJ, Yang JH, Walker NM, Hypponen E, Dunger DB, Ramos-Lopez E, Badenhoop K, Nejentsev S, Todd JA: Association of the vitamin D metabolism gene CYP27B1 with type 1 diabetes. *Diabetes* 2007;56:2616-2621
48. Mori T, Iwasaki Y, Seki Y, Iseki M, Katayama H, Yamamoto K, Takatsu K, Takaki S: Lnk/Sh2b3 controls the production and function of dendritic cells and regulates the induction of IFN-gamma-producing T cells. *J Immunol* 2014;193:1728-1736
49. Steck AK, Xu P, Geyer S, Redondo MJ, Antinozzi P, Wentworth JM, Sosenko J, Onengut-Gumuscu S, Chen WM, Rich SS, Pugliese A, Type 1 Diabetes TrialNet Study G: Can Non-HLA Single Nucleotide Polymorphisms Help Stratify Risk in TrialNet Relatives at Risk for Type 1 Diabetes? *The Journal of clinical endocrinology and metabolism* 2017;102:2873-2880
50. Preuss I, Ludwig MG, Baumgarten B, Bassilana F, Gessier F, Seuwen K, Sailer AW: Transcriptional regulation and functional characterization of the oxysterol/EBI2 system in primary human macrophages. *Biochem Biophys Res Commun* 2014;446:663-668
51. Lu E, Dang EV, McDonald JG, Cyster JG: Distinct oxysterol requirements for positioning naive and activated dendritic cells in the spleen. *Sci Immunol* 2017;2
52. Chen MT, Dong L, Zhang XH, Yin XL, Ning HM, Shen C, Su R, Li F, Song L, Ma YN, Wang F, Zhao HL, Yu J, Zhang JW: ZFP36L1 promotes monocyte/macrophage differentiation by repressing CDK6. *Sci Rep* 2015;5:16229
53. Cooper JD, Smyth DJ, Smiles AM, Plagnol V, Walker NM, Allen JE, Downes K, Barrett JC, Healy BC, Mychaleckyj JC, Warram JH, Todd JA: Meta-analysis of genome-wide association study data identifies additional type 1 diabetes risk loci. *Nature genetics* 2008;40:1399-1401
54. Yoshida H, Hunter CA: The immunobiology of interleukin-27. *Annu Rev Immunol* 2015;33:417-443
55. Do J, Kim D, Kim S, Valentin-Torres A, Dvorina N, Jang E, Nagarajavel V, DeSilva TM, Li X, Ting AH, Vignali DAA, Stohlman SA, Baldwin WM, 3rd, Min B: Treg-specific IL-27Ralpha deletion uncovers a key role for IL-27 in Treg function to control autoimmunity. *Proc Natl Acad Sci U S A* 2017;114:10190-10195
56. Hakonarson H, Grant SF, Bradfield JP, Marchand L, Kim CE, Glessner JT, Grabs R, Casalunovo T, Taback SP, Frackelton EC, Lawson ML, Robinson LJ, Skraban R, Lu Y, Chiavacci RM, Stanley CA, Kirsch SE, Rappaport EF, Orange JS, Monos DS, Devoto M, Qu HQ, Polychronakos C: A genome-wide association study identifies KIAA0350 as a type 1 diabetes gene. *Nature* 2007;448:591-594
57. Tam RC, Li MW, Gao YP, Pang YT, Yan S, Ge W, Lau CS, Chan VS: Human CLEC16A regulates autophagy through modulating mTOR activity. *Exp Cell Res* 2017;352:304-312
58. Schuster C, Gerold KD, Schober K, Probst L, Boerner K, Kim MJ, Ruckdeschel A, Serwold T, Kissler S: The Autoimmunity-Associated Gene CLEC16A Modulates Thymic Epithelial Cell Autophagy and Alters T Cell Selection. *Immunity* 2015;42:942-952
59. Katakam AK, Brightbill H, Franci C, Kung C, Nunez V, Jones C, 3rd, Peng I, Jeet S, Wu LC, Mellman I, Delamarre L, Austin CD: Dendritic cells require NIK for CD40-dependent cross-priming of CD8+ T cells. *Proc Natl Acad Sci U S A* 2015;112:14664-14669
60. Simoncic PD, Bourdeau A, Lee-Loy A, Rohrschneider LR, Tremblay ML, Stanley ER, McGlade CJ: T-cell protein tyrosine phosphatase (Tcptp) is a negative regulator of colony-stimulating factor 1 signaling and macrophage differentiation. *Mol Cell Biol* 2006;26:4149-4160
61. Tahara-Hanaka S, Shibuya K, Kai H, Miyamoto A, Morikawa Y, Ohkochi N, Honda S, Shibuya A: Tumor rejection by the poliovirus receptor family ligands of the DNAM-1 (CD226) receptor. *Blood* 2006;107:1491-1496

## SUPPLEMENTARY DATA

62. Savina A, Peres A, Cebrian I, Carmo N, Moita C, Hacohen N, Moita LF, Amigorena S: The small GTPase Rac2 controls phagosomal alkalinization and antigen crosspresentation selectively in CD8(+) dendritic cells. *Immunity* 2009;30:544-555
63. Kim MJ, Lee W, Park EJ, Park SY: C1qTNF-related protein-6 increases the expression of interleukin-10 in macrophages. *Mol Cells* 2010;30:59-64
64. Cooper JD, Walker NM, Smyth DJ, Downes K, Healy BC, Todd JA, Type IDGC: Follow-up of 1715 SNPs from the Wellcome Trust Case Control Consortium genome-wide association study in type I diabetes families. *Genes Immun* 2009;10 Suppl 1:S85-94
65. Bourgin C, Bourette RP, Arnaud S, Liu Y, Rohrschneider LR, Mouchiroud G: Induced expression and association of the Mona/Gads adapter and Gab3 scaffolding protein during monocyte/macrophage differentiation. *Mol Cell Biol* 2002;22:3744-3756
66. Hamilton-Williams EE, Serreze DV, Charlton B, Johnson EA, Marron MP, Mullbacher A, Slattery RM: Transgenic rescue implicates beta2-microglobulin as a diabetes susceptibility gene in nonobese diabetic (NOD) mice. *Proc Natl Acad Sci U S A* 2001;98:11533-11538
67. Perarnau B, Siegrist CA, Gillet A, Vincent C, Kimura S, Lemonnier FA: Beta 2-microglobulin restriction of antigen presentation. *Nature* 1990;346:751-754
68. Bouloc A, Bagot M, Delaire S, Bensussan A, Boumsell L: Triggering CD101 molecule on human cutaneous dendritic cells inhibits T cell proliferation via IL-10 production. *Eur J Immunol* 2000;30:3132-3139
69. Westwell-Roper CY, Ehses JA, Verchere CB: Resident macrophages mediate islet amyloid polypeptide-induced islet IL-1beta production and beta-cell dysfunction. *Diabetes* 2014;63:1698-1711
70. Wong AS, Mortin-Toth S, Sung M, Canty AJ, Gulban O, Greaves DR, Danska JS: Polymorphism in the innate immune receptor SIRPalpha controls CD47 binding and autoimmunity in the nonobese diabetic mouse. *J Immunol* 2014;193:4833-4844
71. Simpson PB, Mistry MS, Maki RA, Yang W, Schwarz DA, Johnson EB, Lio FM, Alleva DG: Cutting edge: diabetes-associated quantitative trait locus, Idd4, is responsible for the IL-12p40 overexpression defect in nonobese diabetic (NOD) mice. *J Immunol* 2003;171:3333-3337
72. Peng R, Bathjat K, Li Y, Clare-Salzler MJ: Defective maturation of myeloid dendritic cell (DC) in NOD mice is controlled by IDD10/17/18. *Ann N Y Acad Sci* 2003;1005:184-186
73. Summers KL, Marleau AM, Mahon JL, McManus R, Hramiak I, Singh B: Reduced IFN-alpha secretion by blood dendritic cells in human diabetes. *Clin Immunol* 2006;121:81-89
74. Peng R, Li Y, Brezner K, Litherland S, Clare-Salzler MJ: Abnormal peripheral blood dendritic cell populations in type 1 diabetes. *Ann N Y Acad Sci* 2003;1005:222-225
75. Meyers AJ, Shah RR, Gottlieb PA, Zipris D: Altered Toll-like receptor signaling pathways in human type 1 diabetes. *J Mol Med (Berl)* 2010;88:1221-1231
76. Alkanani AK, Rewers M, Dong F, Waugh K, Gottlieb PA, Zipris D: Dysregulated Toll-like receptor-induced interleukin-1beta and interleukin-6 responses in subjects at risk for the development of type 1 diabetes. *Diabetes* 2012;61:2525-2533
77. Xia CQ, Peng R, Chernatynskaya AV, Yuan L, Carter C, Valentine J, Sobel E, Atkinson MA, Clare-Salzler MJ: Increased IFN-alpha-producing plasmacytoid dendritic cells (pDCs) in human Th1-mediated type 1 diabetes: pDCs augment Th1 responses through IFN-alpha production. *J Immunol* 2014;193:1024-1034
78. Nieminen JK, Vakkila J, Salo HM, Ekstrom N, Harkonen T, Ilonen J, Knip M, Vaarala O: Altered phenotype of peripheral blood dendritic cells in pediatric type 1 diabetes. *Diabetes Care* 2012;35:2303-2310
79. Vuckovic S, Withers G, Harris M, Khalil D, Gardiner D, Flesch I, Tepes S, Greer R, Cowley D, Cotterill A, Hart DN: Decreased blood dendritic cell counts in type 1 diabetic children. *Clin Immunol* 2007;123:281-288

## SUPPLEMENTARY DATA

80. Hinkmann C, Knerr I, Hahn EG, Lohmann T, Seifarth CC: Reduced frequency of peripheral plasmacytoid dendritic cells in type 1 diabetes. *Horm Metab Res* 2008;40:767-771
81. Jansen A, van Hagen M, Drexhage HA: Defective maturation and function of antigen-presenting cells in type 1 diabetes. *Lancet* 1995;345:491-492
82. Takahashi K, Honeyman MC, Harrison LC: Impaired yield, phenotype, and function of monocyte-derived dendritic cells in humans at risk for insulin-dependent diabetes. *J Immunol* 1998;161:2629-2635
83. Skarsvik S, Tiittanen M, Lindstrom A, Casas R, Ludvigsson J, Vaarala O: Poor in vitro maturation and pro-inflammatory cytokine response of dendritic cells in children at genetic risk of type 1 diabetes. *Scandinavian journal of immunology* 2004;60:647-652
84. Vasquez AC, Feili-Hariri M, Tan RJ, Morel PA: Qualitative and quantitative abnormalities in splenic dendritic cell populations in NOD mice. *Clin Exp Immunol* 2004;135:209-218
85. Marleau AM, Summers KL, Singh B: Differential contributions of APC subsets to T cell activation in nonobese diabetic mice. *J Immunol* 2008;180:5235-5249
86. O'Keeffe M, Brodnicki TC, Fancke B, Vremec D, Morahan G, Maraskovsky E, Steptoe R, Harrison LC, Shortman K: Fms-like tyrosine kinase 3 ligand administration overcomes a genetically determined dendritic cell deficiency in NOD mice and protects against diabetes development. *Int Immunopharmacol* 2005;17:307-314
87. Prasad SJ, Goodnow CC: Cell-intrinsic effects of non-MHC NOD genes on dendritic cell generation in vivo. *Int Immunopharmacol* 2002;14:677-684
88. Kreuzer D, Nikoopour E, Au BC, Krougly O, Lee-Chan E, Summers KL, Haeryfar SM, Singh B: Reduced interferon-alpha production by dendritic cells in type 1 diabetes does not impair immunity to influenza virus. *Clin Exp Immunol* 2015;179:245-255
89. Boudaly S, Morin J, Berthier R, Marche P, Boitard C: Altered dendritic cells (DC) might be responsible for regulatory T cell imbalance and autoimmunity in nonobese diabetic (NOD) mice. *European cytokine network* 2002;13:29-37
90. Katz JD, Ondr JK, Opoka RJ, Garcia Z, Janssen EM: Cutting edge: merocytic dendritic cells break T cell tolerance to beta cell antigens in nonobese diabetic mouse diabetes. *J Immunol* 2010;185:1999-2003
91. Liang X, Ma L, Thai NL, Fung JJ, Qian S, Lu L: The role of liver-derived regulatory dendritic cells in prevention of type 1 diabetes. *Immunology* 2007;120:251-260
92. Morin J, Chimenes A, Boitard C, Berthier R, Boudaly S: Granulocyte-dendritic cell unbalance in the non-obese diabetic mice. *Cellular immunology* 2003;223:13-25
93. Lee M, Kim AY, Kang Y: Defects in the differentiation and function of bone marrow-derived dendritic cells in non-obese diabetic mice. *J Korean Med Sci* 2000;15:217-223
94. Morel PA, Vasquez AC, Feili-Hariri M: Immunobiology of DC in NOD mice. *J Leukoc Biol* 1999;66:276-280
95. Feili-Hariri M, Morel PA: Phenotypic and functional characteristics of BM-derived DC from NOD and non-diabetes-prone strains. *Clin Immunol* 2001;98:133-142
96. Nikolic T, Bunk M, Drexhage HA, Leenen PJ: Bone marrow precursors of nonobese diabetic mice develop into defective macrophage-like dendritic cells in vitro. *J Immunol* 2004;173:4342-4351
97. Prasad SJ, Goodnow CC: Intrinsic in vitro abnormalities in dendritic cell generation caused by non-MHC non-obese diabetic genes. *Immunology and cell biology* 2002;80:198-206
98. Steptoe RJ, Ritchie JM, Harrison LC: Increased generation of dendritic cells from myeloid progenitors in autoimmune-prone nonobese diabetic mice. *J Immunol* 2002;168:5032-5041
99. Huang Y, Fugier-Vivier IJ, Miller T, Elliott MJ, Xu H, Bozulic LD, Chilton PM, Ildstad ST: Plasmacytoid precursor dendritic cells from NOD mice exhibit impaired function: are they a component of diabetes pathogenesis? *Diabetes* 2008;57:2360-2370
100. Lepault F, Gagnerault MC, Dardenne M, Bach JF: Hematopoietic stem cells and myeloid precursor cells in nonobese diabetic (NOD) mice. *Int J Cell Cloning* 1992;10:173-181

## SUPPLEMENTARY DATA

101. Langmuir PB, Bridgett MM, Bothwell AL, Crispe IN: Bone marrow abnormalities in the non-obese diabetic mouse. *Int Immunol* 1993;5:169-177
102. Welzen-Coppens JM, van Helden-Meeuwesen CG, Drexhage HA, Versnel MA: Abnormalities of dendritic cell precursors in the pancreas of the NOD mouse model of diabetes. *Eur J Immunol* 2012;42:186-194
103. Litherland SA, Grebe KM, Belkin NS, Paek E, Elf J, Atkinson M, Morel L, Clare-Salzler MJ, McDuffie M: Nonobese diabetic mouse congenic analysis reveals chromosome 11 locus contributing to diabetes susceptibility, macrophage STAT5 dysfunction, and granulocyte-macrophage colony-stimulating factor overproduction. *J Immunol* 2005;175:4561-4565
104. Litherland SA, Xie TX, Grebe KM, Davoodi-Semiroomi A, Elf J, Belkin NS, Moldawer LL, Clare-Salzler MJ: Signal transduction activator of transcription 5 (STAT5) dysfunction in autoimmune monocytes and macrophages. *J Autoimmun* 2005;24:297-310
105. Garrigan E, Belkin NS, Alexander JJ, Han Z, Seydel F, Carter J, Atkinson M, Wasserfall C, Clare-Salzler MJ, Amick MA, Litherland SA: Persistent STAT5 phosphorylation and epigenetic dysregulation of GM-CSF and PGS2/COX2 expression in Type 1 diabetic human monocytes. *PloS one* 2013;8:e76919
106. Serreze DV, Gaskins HR, Leiter EH: Defects in the differentiation and function of antigen presenting cells in NOD/Lt mice. *J Immunol* 1993;150:2534-2543
107. Derbinski J, Schulte A, Kyewski B, Klein L: Promiscuous gene expression in medullary thymic epithelial cells mirrors the peripheral self. *Nat Immunol* 2001;2:1032-1039
108. Anderson M, Venanzi E, Klein L, Chen Z, Berzins S, Turley S, von Boehmer H, Bronson R, Dierich A, Benoist C, Mathis D: Projection of an immunological self shadow within the thymus by the aire protein. *Science* 2002;298:1395-1401
109. Gotter J, Brors B, Hergenhahn M, Kyewski B: Medullary epithelial cells of the human thymus express a highly diverse selection of tissue-specific genes colocalized in chromosomal clusters. *J Exp Med* 2004;199:155-166
110. Yip L, Su L, Sheng D, Chang P, Atkinson M, Czesak M, Albert PR, Collier AR, Turley SJ, Fathman CG, Creusot RJ: Deaf1 isoforms control the expression of genes encoding peripheral tissue antigens in the pancreatic lymph nodes during type 1 diabetes. *Nat Immunol* 2009;10:1026-1033
111. Cohen JN, Guidi CJ, Tewalt EF, Qiao H, Rouhani SJ, Ruddell A, Farr AG, Tung KS, Engelhard VH: Lymph node-resident lymphatic endothelial cells mediate peripheral tolerance via Aire-independent direct antigen presentation. *J Exp Med* 2010;207:681-688
112. Yip L, Creusot RJ, Pager CT, Sarnow P, Fathman CG: Reduced DEAF1 function during type 1 diabetes inhibits translation in lymph node stromal cells by suppressing Eif4g3. *Journal of molecular cell biology* 2013;5:99-110
113. Thayer TC, Pearson JA, De Leenheer E, Hanna SJ, Boldison J, Davies J, Tsui A, Ahmed S, Easton P, Siew LK, Wen L, Wong FS: Peripheral Proinsulin Expression Controls Low-Avidity Proinsulin-Reactive CD8 T Cells in Type 1 Diabetes. *Diabetes* 2016;65:3429-3439
114. Narendran P, Neale AM, Lee BH, Ngui K, Steptoe RJ, Morahan G, Madsen O, Dromey JA, Jensen KP, Harrison LC: Proinsulin is encoded by an RNA splice variant in human blood myeloid cells. *Proc Natl Acad Sci U S A* 2006;103:16430-16435
115. Garcia C, Prabakar K, Diez J, Cao Z, Allende G, Zeller M, Dogra R, Mendez A, Rosenkranz E, Dahl U, Ricordi C, Hanahan D, Pugliese A: Dendritic cells in human thymus and periphery display a proinsulin epitope in a transcription-dependent, capture-independent fashion. *J Immunol* 2005;175:2111-2122
116. Grupillo M, Gualtierotti G, He J, Sisino G, Bottino R, Rudert WA, Trucco M, Fan Y: Essential roles of insulin expression in Aire<sup>+</sup> tolerogenic dendritic cells in maintaining peripheral self-tolerance of islet beta-cells. *Cellular immunology* 2012;273:115-123

## SUPPLEMENTARY DATA

117. Sambataro M, Sambado L, Trevisiol E, Cacciatore M, Furlan A, Stefani PM, Segnfreddo E, Durante E, Conte S, Della Bella S, Paccagnella A, Dei Tos AP: Proinsulin-expressing dendritic cells in type 2 neuropathic diabetic patients with and without foot lesions. *FASEB J* 2018;fj201701279RR
118. Poliani PL, Kisand K, Marrella V, Ravanini M, Notarangelo LD, Villa A, Peterson P, Facchetti F: Human peripheral lymphoid tissues contain autoimmune regulator-expressing dendritic cells. *Am J Pathol* 2010;176:1104-1112
119. Kojima H, Fujimiya M, Matsumura K, Nakahara T, Hara M, Chan L: Extrapancreatic insulin-producing cells in multiple organs in diabetes. *Proc Natl Acad Sci U S A* 2004;101:2458-2463
120. Diez J, Park Y, Zeller M, Brown D, Garza D, Ricordi C, Hutton J, Eisenbarth GS, Pugliese A: Differential splicing of the IA-2 mRNA in pancreas and lymphoid organs as a permissive genetic mechanism for autoimmunity against the IA-2 type 1 diabetes autoantigen. *Diabetes* 2001;50:895-900
121. Dogra RS, Vaidyanathan P, Prabakar KR, Marshall KE, Hutton JC, Pugliese A: Alternative splicing of G6PC2, the gene coding for the islet-specific glucose-6-phosphatase catalytic subunit-related protein (IGRP), results in differential expression in human thymus and spleen compared with pancreas. *Diabetologia* 2006;49:953-957
122. de Jong VM, Abreu JR, Verrijn Stuart AA, van der Slik AR, Verhaeghen K, Engelse MA, Blom B, Staal FJ, Gorus FK, Roep BO: Alternative splicing and differential expression of the islet autoantigen IGRP between pancreas and thymus contributes to immunogenicity of pancreatic islets but not diabetogenicity in humans. *Diabetologia* 2013;56:2651-2658
123. Lee J, Epardaud M, Sun J, Becker J, Cheng A, Yonekura A, Heath J, Turley S: Peripheral antigen display by lymph node stroma promotes T cell tolerance to intestinal self. *Nat Immunol* 2007;8:181-190
124. Man SM, Kanneganti TD: Regulation of lysosomal dynamics and autophagy by CTSB/cathepsin B. *Autophagy* 2016;12:2504-2505
125. van Kasteren SI, Overkleef HS: Endo-lysosomal proteases in antigen presentation. *Curr Opin Chem Biol* 2014;23:8-15
126. Floyer T, Brorsson C, Nielsen LB, Miani M, Bang-Bertelsen CH, Friedrichsen M, Overgaard AJ, Berchtold LA, Wiberg A, Poulsen P, Hansen L, Rosinger S, Boehm BO, Ram R, Nguyen Q, Mehta M, Morahan G, Concannon P, Bergholdt R, Nielsen JH, Reinheckel T, von Herrath M, Vaag A, Eizirik DL, Mortensen HB, Storling J, Pociot F: CTSH regulates beta-cell function and disease progression in newly diagnosed type 1 diabetes patients. *Proc Natl Acad Sci U S A* 2014;111:10305-10310
127. Bhatnagar A, Milburn PJ, Lobigs M, Blanden RV, Gautam AM: Nonobese diabetic mice display elevated levels of class II-associated invariant chain peptide associated with I-Ag7 on the cell surface. *J Immunol* 2001;166:4490-4497
128. Silva DG, Socha L, Correcha M, Petrovsky N: Elevated lymphocyte expression of CLIP is associated with type 1 diabetes and may be a useful marker of autoimmune susceptibility. *Ann N Y Acad Sci* 2004;1037:65-68
129. Yan G, Shi L, Penfornis A, Faustman DL: Impaired processing and presentation by MHC class II proteins in human diabetic cells. *J Immunol* 2003;170:620-627
130. Busch R, De Riva A, Hadjinicolaou AV, Jiang W, Hou T, Mellins ED: On the perils of poor editing: regulation of peptide loading by HLA-DQ and H2-A molecules associated with celiac disease and type 1 diabetes. *Expert Rev Mol Med* 2012;14:e15
131. Zhou Z, Reyes-Vargas E, Escobar H, Rudd B, Rockwood AL, Delgado JC, He X, Jensen PE: Type 1 diabetes associated HLA-DQ2 and DQ8 molecules are relatively resistant to HLA-DM mediated release of invariant chain-derived CLIP peptides. *Eur J Immunol* 2016;46:834-845
132. Delong T, Wiles TA, Baker RL, Bradley B, Barbour G, Reisdorph R, Armstrong M, Powell RL, Reisdorph N, Kumar N, Elso CM, DeNicola M, Bottino R, Powers AC, Harlan DM, Kent SC, Mannerling SI, Haskins K: Pathogenic CD4 T cells in type 1 diabetes recognize epitopes formed by peptide fusion. *Science* 2016;351:711-714

## SUPPLEMENTARY DATA

133. Babon JA, DeNicola ME, Blodgett DM, Crevecoeur I, Buttrick TS, Maehr R, Bottino R, Naji A, Kaddis J, Elyaman W, James EA, Haliyur R, Brissova M, Overbergh L, Mathieu C, Delong T, Haskins K, Pugliese A, Campbell-Thompson M, Mathews C, Atkinson MA, Powers AC, Harlan DM, Kent SC: Analysis of self-antigen specificity of islet-infiltrating T cells from human donors with type 1 diabetes. *Nat Med* 2016;
134. Strid J, Lopes L, Marcinkiewicz J, Petrovska L, Nowak B, Chain BM, Lund T: A defect in bone marrow derived dendritic cell maturation in the nonobese diabetic mouse. *Clin Exp Immunol* 2001;123:375-381
135. Sen P, Bhattacharyya S, Wallet M, Wong CP, Poligone B, Sen M, Baldwin AS, Jr., Tisch R: NF-kappa B hyperactivation has differential effects on the APC function of nonobese diabetic mouse macrophages. *J Immunol* 2003;170:1770-1780
136. Price JD, Beauchamp NM, Rahir G, Zhao Y, Rieger CC, Lau-Kilby AW, Tarbell KV: CD8+ dendritic cell-mediated tolerance of autoreactive CD4+ T cells is deficient in NOD mice and can be corrected by blocking CD40L. *J Leukoc Biol* 2014;95:325-336
137. Rahman MJ, Rahir G, Dong MB, Zhao Y, Rodrigues KB, Hotta-Iwamura C, Chen Y, Guerrero A, Tarbell KV: Despite Increased Type 1 IFN, Autoimmune Nonobese Diabetic Mice Display Impaired Dendritic Cell Response to CpG and Decreased Nuclear Localization of IFN-Activated STAT1. *J Immunol* 2016;196:2031-2040
138. Beumer W, Welzen-Coppens JM, van Helden-Meeuwsen CG, Gibney SM, Drechsler HA, Versnel MA: The gene expression profile of CD11c+ CD8alpha- dendritic cells in the pre-diabetic pancreas of the NOD mouse. *PloS one* 2014;9:e103404
139. Poligone B, Weaver DJ, Jr., Sen P, Baldwin AS, Jr., Tisch R: Elevated NF-kappaB activation in nonobese diabetic mouse dendritic cells results in enhanced APC function. *J Immunol* 2002;168:188-196
140. Weaver DJ, Jr., Poligone B, Bui T, Abdel-Motal UM, Baldwin AS, Jr., Tisch R: Dendritic cells from nonobese diabetic mice exhibit a defect in NF-kappa B regulation due to a hyperactive I kappa B kinase. *J Immunol* 2001;167:1461-1468
141. Marleau AM, Singh B: Myeloid dendritic cells in non-obese diabetic mice have elevated costimulatory and T helper-1-inducing abilities. *J Autoimmun* 2002;19:23-35
142. Wheat W, Kupfer R, Gutches DG, Rayat GR, Beilke J, Scheinman RI, Wegmann DR: Increased NF-kappa B activity in B cells and bone marrow-derived dendritic cells from NOD mice. *Eur J Immunol* 2004;34:1395-1404
143. Takahashi K, Satoh J, Oka Y: Lowered expressions of the NF-kappaB family members in dendritic cells from NOD mice are associated with a reduced expression of GATA-2. *Ann N Y Acad Sci* 2008;1150:59-60
144. Angelini F, Del Duca E, Piccinini S, Pacciani V, Rossi P, Manca Bitti ML: Altered phenotype and function of dendritic cells in children with type 1 diabetes. *Clin Exp Immunol* 2005;142:341-346
145. Mollah ZU, Pai S, Moore C, O'Sullivan BJ, Harrison MJ, Peng J, Phillips K, Prins JB, Cardinal J, Thomas R: Abnormal NF-kappa B function characterizes human type 1 diabetes dendritic cells and monocytes. *J Immunol* 2008;180:3166-3175
146. Pruul K, Kisand K, Alnek K, Metskula K, Reimand K, Heilman K, Peet A, Varik K, Peetsalu M, Einberg U, Tillmann V, Uibo R: Differences in B7 and CD28 family gene expression in the peripheral blood between newly diagnosed young-onset and adult-onset type 1 diabetes patients. *Mol Cell Endocrinol* 2015;412:265-271
147. Spatz M, Eibl N, Hink S, Wolf HM, Fischer GF, Mayr WR, Schernthaner G, Eibl MM: Impaired primary immune response in type-1 diabetes. Functional impairment at the level of APCs and T-cells. *Cellular immunology* 2003;221:15-26
148. Zacher T, Knerr I, Rascher W, Kalden JR, Wassmuth R: Characterization of monocyte-derived dendritic cells in recent-onset diabetes mellitus type 1. *Clin Immunol* 2002;105:17-24

## SUPPLEMENTARY DATA

149. Eibl N, Spatz M, Fischer GF, Mayr WR, Samstag A, Wolf HM, Schernthaner G, Eibl MM: Impaired primary immune response in type-1 diabetes: results from a controlled vaccination study. *Clin Immunol* 2002;103:249-259
150. Liu J, Beller D: Aberrant production of IL-12 by macrophages from several autoimmune-prone mouse strains is characterized by intrinsic and unique patterns of NF-kappa B expression and binding to the IL-12 p40 promoter. *J Immunol* 2002;169:581-586
151. Welzen-Coppens JM, van Helden-Meeuwesen CG, Leenen PJ, Drexhage HA, Versnel MA: Reduced numbers of dendritic cells with a tolerogenic phenotype in the prediabetic pancreas of NOD mice. *J Leukoc Biol* 2012;92:1207-1213
152. Alleva DG, Pavlovich RP, Grant C, Kaser SB, Beller DI: Aberrant macrophage cytokine production is a conserved feature among autoimmune-prone mouse strains: elevated interleukin (IL)-12 and an imbalance in tumor necrosis factor-alpha and IL-10 define a unique cytokine profile in macrophages from young nonobese diabetic mice. *Diabetes* 2000;49:1106-1115
153. Lee MS, Kwon HJ, Kim HS: Macrophages from nonobese diabetic mouse have a selective defect in IFN-gamma but not IFN-alpha/beta receptor pathway. *J Clin Immunol* 2012;32:753-761
154. Giulietti A, Stoffels K, Decallonne B, Overbergh L, Mathieu C: Monocytic expression behavior of cytokines in diabetic patients upon inflammatory stimulation. *Ann N Y Acad Sci* 2004;1037:74-78
155. Thompson WS, Pekalski ML, Simons HZ, Smyth DJ, Castro-Dopico X, Guo H, Guy C, Dunger DB, Arif S, Peakman M, Wallace C, Wicker LS, Todd JA, Ferreira RC: Multi-parametric flow cytometric and genetic investigation of the peripheral B cell compartment in human type 1 diabetes. *Clin Exp Immunol* 2014;177:571-585
156. Deng C, Xiang Y, Tan T, Ren Z, Cao C, Huang G, Wen L, Zhou Z: Altered Peripheral B-Lymphocyte Subsets in Type 1 Diabetes and Latent Autoimmune Diabetes in Adults. *Diabetes care* 2016;39:434-440
157. Jagannathan M, McDonnell M, Liang Y, Hasturk H, Hetzel J, Rubin D, Kantarci A, Van Dyke TE, Ganley-Leal LM, Nikolajczyk BS: Toll-like receptors regulate B cell cytokine production in patients with diabetes. *Diabetologia* 2010;53:1461-1471
158. Devaraj S, Dasu MR, Rockwood J, Winter W, Griffen SC, Jialal I: Increased toll-like receptor (TLR) 2 and TLR4 expression in monocytes from patients with type 1 diabetes: further evidence of a proinflammatory state. *The Journal of clinical endocrinology and metabolism* 2008;93:578-583
159. Plesner A, Greenbaum CJ, Gaur LK, Ernst RK, Lernmark A: Macrophages from high-risk HLA-DQB1\*0201/\*0302 type 1 diabetes mellitus patients are hypersensitive to lipopolysaccharide stimulation. *Scandinavian journal of immunology* 2002;56:522-529
160. Kayserova J, Vcelakova J, Stechova K, Dudkova E, Hromadkova H, Sumnik Z, Kolouskova S, Spisek R, Sediva A: Decreased dendritic cell numbers but increased TLR9-mediated interferon-alpha production in first degree relatives of type 1 diabetes patients. *Clin Immunol* 2014;153:49-55
161. Gomez-Tourino I, Sanchez-Espinel C, Hernandez-Fernandez A, Gonzalez-Fernandez A, Pena-Gonzalez E, Rodriguez J, Garcia-Lopez JM, Varela-Calvino R: Galectin-1 synthesis in type 1 diabetes by different immune cell types: reduced synthesis by monocytes and Th1 cells. *Cellular immunology* 2011;271:319-328
162. Munn DH, Mellor AL: Indoleamine 2,3 dioxygenase and metabolic control of immune responses. *Trends Immunol* 2013;34:137-143
163. Saxena V, Ondr JK, Magnussen AF, Munn DH, Katz JD: The countervailing actions of myeloid and plasmacytoid dendritic cells control autoimmune diabetes in the nonobese diabetic mouse. *J Immunol* 2007;179:5041-5053
164. Grohmann U, Fallarino F, Bianchi R, Orabona C, Vacca C, Fioretti MC, Puccetti P: A defect in tryptophan catabolism impairs tolerance in nonobese diabetic mice. *J Exp Med* 2003;198:153-160

## SUPPLEMENTARY DATA

165. Ueno A, Cho S, Cheng L, Wang J, Hou S, Nakano H, Santamaria P, Yang Y: Transient upregulation of indoleamine 2,3-dioxygenase in dendritic cells by human chorionic gonadotropin downregulates autoimmune diabetes. *Diabetes* 2007;56:1686-1693
166. Hosseini-Tabatabaei A, Jalili RB, Li Y, Kilani RT, Moeen Rezakhanlou A, Ghahary A: Mechanism underlying defective interferon gamma-induced IDO expression in non-obese diabetic mouse fibroblasts. *PloS one* 2012;7:e37747
167. Bscheider M, Butcher EC: Vitamin D immunoregulation through dendritic cells. *Immunology* 2016;148:227-236
168. Shahijanian F, Parnell GP, McKay FC, Gatt PN, Shojoei M, O'Connor KS, Schibeci SD, Brilot F, Liddle C, Batten M, Consortium ANMSG, Stewart GJ, Booth DR: The CYP27B1 variant associated with an increased risk of autoimmune disease is underexpressed in tolerizing dendritic cells. *Hum Mol Genet* 2014;23:1425-1434
169. Ramos-Lopez E, Bruck P, Jansen T, Pfeilschifter JM, Radeke HH, Badenhoop K: CYP2R1-, CYP27B1- and CYP24-mRNA expression in German type 1 diabetes patients. *J Steroid Biochem Mol Biol* 2007;103:807-810
170. Rose K, Penna-Martinez M, Klahold E, Karger D, Shoghi F, Kahles H, Bayer M, Hintermann E, Pfeilschifter JM, Badenhoop K, Ramos-Lopez E, Christen U: Influence of the vitamin D plasma level and vitamin D-related genetic polymorphisms on the immune status of patients with type 1 diabetes: a pilot study. *Clin Exp Immunol* 2013;171:171-185
171. Paronen J, Klemetti P, Kantele JM, Savilahti E, Perheentupa J, Akerblom HK, Vaarala O: Glutamate decarboxylase-reactive peripheral blood lymphocytes from patients with IDDM express gut-specific homing receptor alpha4beta7-integrin. *Diabetes* 1997;46:583-588
172. Hanninen A, Salmi M, Simell O, Andrew D, Jalkanen S: Recirculation and homing of lymphocyte subsets: dual homing specificity of beta 7-integrin(high)-lymphocytes in nonobese diabetic mice. *Blood* 1996;88:934-944
173. Hanninen A, Salmi M, Simell O, Jalkanen S: Mucosa-associated (beta 7-integrinhigh) lymphocytes accumulate early in the pancreas of NOD mice and show aberrant recirculation behavior. *Diabetes* 1996;45:1173-1180
174. Bouma G, Coppens JM, Mourits S, Nikolic T, Sozzani S, Drexhage HA, Versnel MA: Evidence for an enhanced adhesion of DC to fibronectin and a role of CCL19 and CCL21 in the accumulation of DC around the pre-diabetic islets in NOD mice. *Eur J Immunol* 2005;35:2386-2396
175. Baev DV, Caielli S, Ronchi F, Coccia M, Facciotti F, Nichols KE, Falcone M: Impaired SLAM-SLAM homotypic interaction between invariant NKT cells and dendritic cells affects differentiation of IL-4/IL-10-secreting NKT2 cells in nonobese diabetic mice. *J Immunol* 2008;181:869-877
176. Beaudoin L, Laloux V, Novak J, Lucas B, Lehuen A: NKT cells inhibit the onset of diabetes by impairing the development of pathogenic T cells specific for pancreatic beta cells. *Immunity* 2002;17:725-736
177. Bouma G, Nikolic T, Coppens JM, van Helden-Meeuwsen CG, Leenen PJ, Drexhage HA, Sozzani S, Versnel MA: NOD mice have a severely impaired ability to recruit leukocytes into sites of inflammation. *Eur J Immunol* 2005;35:225-235
178. Kriegel MA, Rathinam C, Flavell RA: Pancreatic islet expression of chemokine CCL2 suppresses autoimmune diabetes via tolerogenic CD11c+ CD11b+ dendritic cells. *Proc Natl Acad Sci U S A* 2012;109:3457-3462
179. Solomon M, Balasa B, Sarvetnick N: CCR2 and CCR5 chemokine receptors differentially influence the development of autoimmune diabetes in the NOD mouse. *Autoimmunity* 2010;43:156-163
180. Hanifi-Moghaddam P, Kappler S, Seissler J, Muller-Scholze S, Martin S, Roep BO, Strassburger K, Kolb H, Schloot NC: Altered chemokine levels in individuals at risk of Type 1 diabetes mellitus. *Diabetic medicine : a journal of the British Diabetic Association* 2006;23:156-163

## SUPPLEMENTARY DATA

181. Carvalho-Pinto C, Garcia MI, Gomez L, Ballesteros A, Zaballos A, Flores JM, Mellado M, Rodriguez-Frade JM, Balomenos D, Martinez AC: Leukocyte attraction through the CCR5 receptor controls progress from insulitis to diabetes in non-obese diabetic mice. *Eur J Immunol* 2004;34:548-557
182. Klementowicz JE, Mahne AE, Spence A, Nguyen V, Satpathy AT, Murphy KM, Tang Q: Cutting Edge: Origins, Recruitment, and Regulation of CD11c<sup>+</sup> Cells in Inflamed Islets of Autoimmune Diabetes Mice. *J Immunol* 2017;199:27-32
183. Dubois-Laforgue D, Hendel H, Caillat-Zucman S, Zagury JF, Winkler C, Boitard C, Timsit J: A common stromal cell-derived factor-1 chemokine gene variant is associated with the early onset of type 1 diabetes. *Diabetes* 2001;50:1211-1213
184. Ide A, Kawasaki E, Abiru N, Sun F, Fukushima T, Takahashi R, Kuwahara H, Fujita N, Kita A, Oshima K, Uotani S, Yamasaki H, Yamaguchi Y, Kawabata Y, Fujisawa T, Ikegami H, Eguchi K: Stromal-cell derived factor-1 chemokine gene variant is associated with type 1 diabetes age at onset in Japanese population. *Human immunology* 2003;64:973-978
185. Welzen-Coppens JM, van Helden-Meeuwsen CG, Leenen PJ, Drexhage HA, Versnel MA: The kinetics of plasmacytoid dendritic cell accumulation in the pancreas of the NOD mouse during the early phases of insulitis. *PloS one* 2013;8:e55071
186. Feili-Hariri M, Dong X, Alber S, Watkins S, Salter R, Morel P: Immunotherapy of NOD mice with bone marrow-derived dendritic cells. *Diabetes* 1999;48:2300-2308
187. Papaccio G, Nicoletti F, Pisanti FA, Bendtzen K, Galdieri M: Prevention of spontaneous autoimmune diabetes in NOD mice by transferring *in vitro* antigen-pulsed syngeneic dendritic cells. *Endocrinology* 2000;141:1500-1505
188. Krueger T, Wohlrab U, Klucken M, Schott M, Seissler J: Autoantigen-specific protection of non-obese diabetic mice from cyclophosphamide-accelerated diabetes by vaccination with dendritic cells. *Diabetologia* 2003;46:1357-1365
189. Lo J, Peng RH, Barker T, Xia CQ, Clare-Salzler MJ: Peptide-pulsed immature dendritic cells reduce response to beta cell target antigens and protect NOD recipients from type I diabetes. *Ann N Y Acad Sci* 2006;1079:153-156
190. Marin-Gallen S, Clemente-Casares X, Planas R, Pujol-Autonell I, Carrascal J, Carrillo J, Ampudia R, Verdaguer J, Pujol-Borrell R, Borras FE, Vives-Pi M: Dendritic cells pulsed with antigen-specific apoptotic bodies prevent experimental type 1 diabetes. *Clin Exp Immunol* 2010;160:207-214
191. Haase C, Yu L, Eisenbarth G, Markholst H: Antigen-dependent immunotherapy of non-obese diabetic mice with immature dendritic cells. *Clin Exp Immunol* 2010;160:331-339
192. Gibson VB, Nikolic T, Pearce VQ, Demengeot J, Roep BO, Peakman M: Proinsulin multi-peptide immunotherapy induces antigen-specific regulatory T cells and limits autoimmunity in a humanized model. *Clin Exp Immunol* 2015;182:251-260
193. Funda DP, Golias J, Hudcovic T, Kozakova H, Spisek R, Palova-Jelinkova L: Antigen Loading (e.g., Glutamic Acid Decarboxylase 65) of Tolerogenic DCs (tolDCs) Reduces Their Capacity to Prevent Diabetes in the Non-Obese Diabetes (NOD)-Severe Combined Immunodeficiency Model of Adoptive Cotransfer of Diabetes As Well As in NOD Mice. *Frontiers in immunology* 2018;9:290